

Chapter 1: Ecological Units of the Highlands

This report is one of a five developed for the Ozark-Ouachita Highlands Assessment. The team that produced this report was comprised of scientists from a variety of disciplines—wildlife biologists, foresters, ecologists, pathologists, and entomologists. They had in common an interest in the terrestrial vegetation and animal species of the Highlands and a willingness to examine the status of these “terrestrial resources.”

Questions about the terrestrial animals, plants, and biological communities of the Highlands were developed at a series of Terrestrial Team meetings in the summer of 1997. No new information was collected in the field to address these questions. Instead, existing data from various agencies and individuals were gathered and analyzed. Two concerns emerged powerfully as sources of information were identified: while an overwhelming amount of information exists for the Highlands, many species have not been studied and the information that is available from various sources is often in different formats, covers different time periods, and/or covers different geographic areas.

One early and continuing challenge was how best to organize and present the information. The Terrestrial Team decided to use the ecologically defined units presented in this chapter to provide a consistent framework throughout the report. Chapter 2 presents an overview of prehistoric and more recent change in the Highlands. Current vegetation cover is examined in some detail in Chapter 3, and silvicultural practices receive in-depth treatment in Chapter 4. Chapter 5 examines many plant and wildlife species of special concern, and Chapter 6 covers current and future biological threats to forest resources.

Question 1.1: What are the ecological boundaries and subsections of the Highlands?

Key Findings

1. The map of sections and subsections of the Ozark-Ouachita Highlands (Keys and others 1995) was revised to better meet the needs of the Assessment. Changes were largely limited to the Arkansas and Oklahoma portions of the Assessment area.
2. The modified map of sections and subsections of the Highlands of Arkansas and Oklahoma is the first such delineation for Oklahoma and provides significant advancements over earlier maps for Arkansas by Croneis (1930) and Foti (1974).

The Ozark-Ouachita Highlands Assessment Area

This report contains data and key findings in the context of an ecologically defined area—the Ozark-Ouachita Highlands (fig. 1.1). Although the term “Interior Highlands” is familiar to geographers and biologists, it is not commonly recognized in other circles. Most people are more familiar with “the Ozarks” and “the Ouachita Mountains” than they are with the older geographic name for the two areas. Thus, the Assessment team leaders chose to use a term likely to be more widely recognized—the Ozark-Ouachita Highlands.

The Ozark-Ouachita Highlands is an appropriate area to assess because it has relatively consistent ecological characteristics and is distinctively different from surrounding landscapes. Consisting mostly of hilly to mountainous topography over substrates of Paleozoic age, the Highlands have long been recognized as a distinct physiographic and natural region (Fenneman 1938, Braun 1950). Upland hardwood and upland pine-hardwood forests characterize much of the area. Lower-lying plains with more recent geological substrates also occur in portions of the Highlands, including much of the Arkansas Valley section. The vegetation of these plains ranges from tallgrass prairie to lowland pine-hardwood and bottomland hardwood forests.

Even though the Ozark-Ouachita Highlands have some consistent general characteristics, there are also striking differences within it. Therefore, most descriptions and studies divide the region into smaller, more uniform areas. Authors have generally recognized at least two provinces, the Ozark Mountains and the Ouachita Mountains (Fenneman 1938, Thornbury 1965, Braun 1950, Croneis 1930, Foti 1974). Sometimes, the Arkansas Valley has been considered a separate province or natural division (Foti 1976, Pell 1983, Omernik 1987). While most authors treated the Boston Mountains as a subdivision of the Ozark Mountains, Omernik (1987) recognized it as a separate ecoregion (natural division). These provinces, natural divisions, or sections are often subdivided as well.

Data Sources

To facilitate agency ecosystem management efforts, the USDA Forest Service developed a new regionalization framework for the Eastern United States (Keys and others 1995, henceforth referred to as “Keys and others” or the “Keys map”) based on a national map of ecoregions of the United States by Bailey and others (1994). The new framework is hierarchical (like older efforts) but is based on a more holistic consideration of landscape properties than some earlier maps, with climate and soil playing prominent roles along with physiography. The new framework is also designed to rationally subdivide landscapes in ways meaningful to ecosystem management. The older and newer maps coincide most closely at the level of section (Keys and others), province (Fenneman 1938), and natural division (Foti 1974). Although differences occur at this level they are usually in the form of one unit in one system equating to two units in another system. The new framework is often more detailed at lower levels in the hierarchy than older maps.

The Terrestrial Team examined the Keys and others (1995) framework to determine whether the ecological units and their boundaries were adequate for Assessment purposes. Important considerations were that the sections and subsections and their boundaries be ecologically meaningful and consistent across State lines. Examination of the Keys map and comparison

with other regional maps and geological and topographical base maps revealed that sections and subsections and their boundaries were not consistently meaningful and accurate across the Assessment area. The Missouri units and their boundaries have been settled for years; therefore, the Keys map simply adopted those boundaries, and changes needed for the Assessment were very minor.

In contrast, the Arkansas units and boundaries needed considerable revision because: (1) the Keys and others (1995) approach departs substantially and without convincing rationale from long-standing delineations (Croneis 1930, Foti 1974) and (2) locally created maps were not available. The Keys map is also problematic in Oklahoma, because in that State only general regions have been defined (OK BTF 1996), boundaries are not detailed, and subdivisions are not mapped. Furthermore, the Keys map appears to be derived from low-detail base maps, and its boundaries were judged to be too general for this Assessment. The Keys map and supporting materials do not explicitly define the source or rationale for boundaries; therefore, revision of the map sometimes required a determination of the defining physical feature and use of an appropriate base map.

Although production of the new map (fig. 1.1) involved many changes to the Keys and others (1995) map, few changes were made in the list of sections and subsections. (A map illustrating the changes the team made to subsection boundaries is available on the Web site for the Assessment, <<http://www.fs.fed.us.oonf>>.) The emphasis was on employing clearly-stated boundary definitions that in most cases were first articulated by Croneis (1930), and then using appropriate digital base maps to create an accurate final product. Many changes were made to the Croneis (1930) and Foti (1974) maps, however, primarily by adding detail to the older maps. (See, for examples, the White River Hills and Central Plateau subsections [which are nested within the Salem Plateau of Croneis 1930], the Upper and Lower Boston Mountains subsections [nested within the Boston Mountain subdivision of Foti 1974], or the three new subsections within the Arkansas Valley.)

Croneis (1930) and Foti (1974) presented rationales for many regional boundaries in Arkansas and Oklahoma, most of which were adopted for this revision. All boundaries are based on either geology or topography,

although soils maps were used for comparison in some cases. The geologic base map was the 1:2,500,000-scale geology of the conterminous United States (Schruben and others 1994). The topographic base map was created for this project from 30-meter (m) USGS digital elevation model files by the Spatial Analysis Laboratory of the School of Forest Resources, University of Arkansas at Monticello.

Ecological Units

The following discussion describes the ecological units used in the Ozark-Ouachita Highlands area and the factors on which the boundaries were based along with changes from the Keys map. Alphanumeric codes used here are the same as those used in the Keys map.

The modified map of sections and subsections of the Highlands of Arkansas and Oklahoma (fig. 1.1) is the first such delineation for Oklahoma and provides four significant advancements over earlier maps for Arkansas by Croneis (1930) and Foti (1974):

- Boundaries are defined and mapped consistently across the three States sharing the Highlands;
- Boundaries based on topography are much more accurate than previous maps due to the use of 30-m digital elevation models;
- Changes in section and subsection definitions that have occurred since production of the earlier maps are incorporated; and
- The map produced by this team is in digital form and freely available on the Assessment Web site.

Ozark Highlands Section (222A)

Six subsections in Missouri and Illinois were not included in the Assessment area: 222Ai (Prairie Ozark Border); 222Aj (Inner Ozark Border); 222Ak (Outer Ozark Border); 222Ao (Mississippi River Alluvial Plain); 222Ap (Missouri River Alluvial Plain); and 222Aq (Illinois Ozarks). These were excluded because they are on the periphery of the region, are not included in some data sets used for the Assessment, and would have complicated the analysis by extending it to an additional State. Following are the subsections that were included in the Assessment area and brief descriptions of each.

222Aa—The St. Francis Knobs and Basins, which cover 1,108,009 acres (ac), occur only in Missouri. They consist of steep hills (but not an eroded plateau like the other Ozark subsections) that are 400 to 1,700 feet (ft) in elevation and underlain by Cambrian and Precambrian igneous and sedimentary rocks. The subsection is covered with acid glades, oak woodlands, and dry-mesic oak forests. No changes were made in the Keys map boundaries.

222Ab—The Central Plateau subsection occurs in Missouri (5,006,390 ac) and Arkansas (1,335,220 ac) and consists of irregular plains 300 to 1,600 ft in elevation with karst features on Ordovician cherty dolomite, sandstone, and cherty clay residuum covered with prairies, oak woodlands, and dry-mesic oak forests. The Keys map boundary with the White River Hills subsection was altered to follow the break in topography between these subsections.

222Ac—The Osage River Hills (1,550,855 ac) occur only in Missouri, where they consist of hills with entrenched valleys 600 to 1,100 ft in elevation that were formed by streams downcutting to the Osage River. Underlain by Ordovician cherty dolomite, sandstone, and cherty clay residuum, this subsection is covered with pine-oak and oak woodlands and forests. No changes were made in the Keys map boundaries.

222Ad—The Gasconade River Hills subsection (1,098,006 ac) occurs only in Missouri. Similar to the Osage River Hills, this subsection consists of hills with entrenched valleys and karst features and was formed by streams downcutting to the Gasconade River. Underlain by Ordovician cherty dolomite, sandstone, and cherty clay residuum, the Gasconade River Hills range from 600 to 1,100 ft in elevation and are covered with pine-oak and oak woodlands and forests. No changes were made in the Keys map boundaries.

222Ae—The Meramac River Hills (1,136,219 ac) occur only in Missouri. (The correct spelling of the river for which this subsection is named is “Meramec,” but the team left the Keys and others [1995] name unchanged.) This subsection consists of hills with entrenched valleys formed by streams downcutting to the Meramec River. Underlain by Cambrian and Ordovician cherty dolomite and cherty clay residuum and ranging from 500 to 1,300 ft in elevation, this subsection is covered with pine-oak and oak woodlands and forests. No changes were made in the Keys map boundaries.

222Af—The Current River Hills (1,563,186 ac) occur in Missouri and consist of entrenched valleys with karst features formed by streams downcutting to the Current River. Underlain by Cambrian and Ordovician cherty dolomite and sandstone with cherty clay residuum and ranging from 400 to 1,300 ft in elevation, the Current River Hills are covered with pine-oak and oak woodlands and forests. No changes were made in the Keys map boundaries.

222Ag—The White River Hills subsection occurs in Missouri (2,155,950 ac) and Arkansas (1,577,221 ac) and consists of hills with entrenched valleys and karst features. It was formed by streams downcutting to the White River. Underlain by Ordovician cherty dolomite with cherty clay residuum and ranging from 600 to 1,600 ft in elevation, this subsection is covered with alkaline glades and oak woodlands and forests. Changes were made in the Arkansas portion of the Keys map boundaries to better follow the break in topography from the surrounding plains.

222Ah—The Elk River Hills occur in Missouri (356,326 ac), Arkansas (57,433 ac), and Oklahoma (32,334 ac) and consist of hills with entrenched valleys and karst features. Formed by streams downcutting to the Neosho River, the subsection is underlain by Mississippian cherty limestone with cherty clay residuum, ranges from 900 to 1,400 ft in elevation, and is covered with oak woodlands and forests. Changes were made in the Arkansas portion of the Keys map boundaries to better follow the break in topography from the surrounding subsections.

222Ai—The Black River Ozark Border (859,059 ac) occurs only in Missouri. It consists of irregular plains and low hills with karst features. Underlain by Ordovician sandstone and cherty dolomite with cherty clay residuum, this subsection ranges from 300 to 900 ft in elevation and is covered with pine-oak and oak woodlands and forests and oak-sweetgum forests. The Keys map boundaries were modified to eliminate this subsection from Arkansas.

222Am—The Springfield Plain lies in Missouri (3,136,051 ac) and Oklahoma (161,881 ac) and is a smooth plain with karst features underlain by Mississippian limestone (sometimes very cherty) and cherty clay residuum. Ranging from 800 to 1,700 ft in elevation, this subsection is covered with prairie and oak

woodlands and forest. No changes were made in the Keys map boundaries.

222An—The Springfield Plateau subsection occurs in Oklahoma (1,486,718 ac), Arkansas (1,579,841 ac), and Missouri (56,326 ac) and consists of smooth to irregular plains 800 to 1,400 ft in elevation with karst features. Underlain by Mississippian limestone (sometimes very cherty) and cherty clay residuum, this subsection is covered with prairie and oak woodlands and forest, alkaline, and acid glades. Detail changes were made in the Keys map boundaries to better follow the break in topography to the Elk River Hills and to more closely follow the boundaries with older and younger geological substrates throughout the rest of the subsection perimeter.

Boston Mountains Section (M222A)

In earlier maps, with the exception of Omernik (1987), this section was treated as a subsection or equivalent.

M222Aa—The Upper Boston Mountains (1,106,642 ac) occur only in Arkansas. They consist of low mountains 1,000 to 2,700 ft in elevation underlain by Pennsylvanian sandstone and shale with sandy residuum and loamy colluvium. This subsection is covered with oak woodlands and forests. Detail changes were made in the Keys map boundaries to better follow the geologic boundary with the Springfield Plateau and to better follow the corresponding land type association boundaries developed by the Ozark-St. Francis National Forests elsewhere along the perimeter of the subsection. This subsection was defined on the basis of elevation (approximating the 1,800-ft elevation contour), which corresponds to areas of lower temperature and higher rainfall and consequent changes in plant community composition. The Keys map name for this subsection (Boston Mountains) and the following subsection (Boston Hills) were changed to reflect that both are parts of the vernacular and physiographic Boston Mountains.

M222Ab—The Lower Boston Mountains subsection occurs in Oklahoma (834,553 ac) and Arkansas (2,471,699 ac) and consists of high hills 500 to 1,800 ft in elevation underlain by Pennsylvanian sandstone and shale with sandy residuum and loamy colluvium. The

Lower Boston Mountains are covered with pine-oak and oak woodlands and forests. Detail changes were made to the Keys map boundaries to better follow the corresponding landtype association boundaries developed by the Ozark-St. Francis National Forests for the Upper Boston Mountains and to better follow the boundary with younger and older geologic substrates elsewhere along the northern, eastern, and western perimeter of the subsection and the topographically defined southern boundary (the escarpment to the Arkansas Valley section [Croneis 1930, Foti 1974]). The Keys map name for this subsection (Boston Hills) was changed as explained in the description of the Upper Boston Mountains.

Arkansas Valley Section (231G)

231Ga—The Eastern Arkansas Valley (1,490,182 ac) lies entirely in Arkansas, where it consists of plains with hills 300 to 500 ft in elevation. Underlain by Pennsylvanian sandstone and shale with sandy residuum, this subsection is covered with pine-oak and pine woodlands and forests. Northern and eastern boundaries were modified in detail to better match topographic and geologic boundaries, respectively. The southern boundary was redefined to match the traditional physiographic boundary, Cadron Ridge (Croneis 1930, Foti 1974). The southwestern boundary was redefined to place all Arkansas River bottomlands within the Western Arkansas Valley subsection; topographic and geologic boundaries also contributed to the modified subsection boundary. The Keys map name was changed to eliminate “and Ridges” since the redefined southern boundary eliminated the most prominent structural ridges from the subsection (one of the reasons for redefining that boundary).

231Gb—The Western Arkansas Valley Mountains occurs in Oklahoma (494,643 ac) and Arkansas (433,498 ac). It consists of low mountains and ridges and some wide valleys as well. Ranging from 750 to 2,800 ft in elevation, the Western Arkansas Valley Mountains are underlain by Pennsylvanian sandstone and shale with sandy residuum and covered with pine-oak and oak woodlands and forests and prairies. The eastern, northern, and western boundaries as delineated on the Keys map were modified somewhat to better

include the mountains and exclude the plains that were continuations of those in the Western Arkansas Valley. The southern boundary was changed to follow the northern boundary of the physiographic Ouachita Mountains (Croneis 1930, Foti 1974). The Keys map name (Mount Magazine) was changed to reflect the importance of other mountains within this subsection.

231Gc—The Western Arkansas Valley subsection includes portions of Oklahoma (829,099 ac) and Arkansas (1,354,977 ac) and consists of plains, low hills, and ridges 300 to 1,000 ft in elevation underlain by Pennsylvanian sandstone and shale with sandy and clayey residuum along with Holocene sandy alluvium. This subsection is covered with pine-oak and oak woodlands and forests, substantial bottomland forests, and prairies. One major low mountain, Petit Jean Mountain, was included within this section because it was disjunct from the Western Arkansas Valley Mountains, in which it would otherwise have been included. The northern, eastern, and southern boundaries of the Keys map were refined based on topography and geology to place all of the Arkansas River alluvial plains, the most extensive alluvial plains of its major tributaries, and almost all of the Pennsylvanian eroded plains within this subsection. A substantial area that extended up the Canadian River at the western end of this subsection on the Keys map was eliminated on the basis of geology, topography, and the definition of the Arkansas Valley as lying between the Ouachita Mountains and the uplifted plateaus of the Ozark Mountains (Croneis 1930).

Ouachita Mountains Section (M231A)

M231Aa—The Fourche Mountains occur in Oklahoma (743,093 ac) and Arkansas (2,148,080 ac) where they form open, low to relatively high mountain ridges, often with wide valleys. Elevations range from 750 to over 2,600 ft, among the highest in the Assessment area. Ridges are underlain by Pennsylvanian and Mississippian sandstone and shale valleys by sandy residuum. Slopes and ridges are covered with pine-oak and oak woodlands and forests. The northern boundary was modified from Keys to coincide with the physiographic boundary based on topography (Croneis 1930, Foti 1974). The southern boundary was modified to match the boundary with Mississippian Arkansas


Novaculite and toward the west to follow the long narrow ridges and include the Pennsylvanian Jackfork Sandstone.

M231Ab—The Western Ouachita Mountains subsection occurs in Oklahoma (1,623,109 ac) and Arkansas (109,249 ac) and consists of open high hills and low mountains, often with wide valleys, with elevations ranging from 750 to 2,500 ft. The subsection is underlain by Mississippian sandstone and shale with clayey colluvium, covered with pine-oak and oak woodlands and forests, along with prairies. The Keys map boundaries were modified using geology (Arkansas Novaculite) to eliminate portions of the Central Ouachita Mountains from this subsection. The word “Central” was eliminated from the Keys map name (West Central Ouachita Mountains) because a substantial part of the subsection lies along the southern boundary of the Ouachita Mountains section.

M231Ac—The Central Ouachita Mountains occur in Oklahoma (244,015 ac) and Arkansas (1,401,574 ac).

They consist of open high hills and low mountains, often with wide valleys, and they range from 750 to 2,500 ft in elevation. The Central Ouachita Mountains are underlain by Mississippian sandstone and shale with clayey colluvium and covered with pine-oak and oak woodlands and forests. The Keys map boundaries were modified using geology (Arkansas Novaculite); a large disjunct area with consistent characteristics is newly delineated in southeastern Oklahoma. The Keys map name was changed by dropping “East” as it was no longer needed (because of the name change to Western Ouachita Mountains).

M231Ad—The Athens Piedmont Plateau occurs in Oklahoma (56,546 ac) and Arkansas (837,602 ac). It consists of open high hills underlain by Mississippian (with small amounts of Pennsylvanian) sandstone and shale with sandy and clay-loam colluvium covered with pine-oak and pine woodlands and forests. The Keys map boundary was refined using geology (Arkansas Novaculite) for north and west boundaries and Tertiary and Cretaceous deposits on the south and east.



Section name	Map code	Subsection name	Subsection number
Ozark Highlands	1	St.Francois Knobs and Basins	222Aa
Ozark Highlands	2	Central Plateau	222Ab
Ozark Highlands	3	Osage River Hills	222Ac
Ozark Highlands	4	Gasconade River Hills	222Ad
Ozark Highlands	5	Meramac River Hills	222Ae
Ozark Highlands	6	Current River Hills	222Af
Ozark Highlands	7	White River Hills	222Ag
Ozark Highlands	8	Elk River Hills	222Ah
Ozark Highlands	9	Black River Ozark Border	222A1
Ozark Highlands	10	Springfield Plain	222Am
Ozark Highlands	11	Springfield Plateau	222An
Boston Mountains	12	Upper Boston Mountains	M222Aa
Boston Mountains	13	Lower Boston Mountains	M222Ab
Arkansas Valley	14	Eastern Arkansas Valley	231Ga
Arkansas Valley	15	Western Arkansas Valley Mountains	231Gb
Arkansas Valley	16	Western Arkansas Valley	231Gc
Ouachita Mountains	17	Fourche Mountains	M231Aa
Ouachita Mountains	18	Western Ouachita Mountains	M231Ab
Ouachita Mountains	19	Central Ouachita Mountains	M231Ac
Ouachita Mountains	20	Athens Piedmont Plateau	M231Ad

Figure 1.1—Ecological sections and subsections of the Ozark-Ouachita Highlands Assessment area (sections and subsections modified from Keys and others 1995).

References

- Abrams, Marc D. 1992. Fire and the development of oak forests. *Bioscience*. 42(5): 346–353.
- Albert, Lois E.; Wyckoff, Don G. 1981. Ferndale bog and natural lake: five thousand years of environmental change in southeastern Oklahoma. No. 7. Oklahoma City, OK: Oklahoma Archeological Survey. 121 p.
- Allard, D.J. 1990. Southeastern United States ecological community classification. Interim report, Version 1.2. Chapel Hill, NC: The Nature Conservancy, Southeast Regional Office. 96 p.
- Ammon, Vernon; Nebeker, T. Evan; Filer, Ted H. [and others]. 1989. Oak decline. Tech. Bull. 161. Mississippi State, MS: Mississippi Agricultural and Forestry Experiment Station. 15 p.
- Anderson, J.F.; Furniss, W.E. 1983. Epidemic of *urticaria* associated with first-instar larvae of the gypsy moth (Lepidoptera: Lymantriidae). *Journal of Medical Entomology*. 20(2): 146–150.
- Anderson, R.L. 1988. The butternut: outlook grim. *American Forests*. 94(9/10): 49.
- Anderson, R.L.; Knighten, J.L.; Windham, M. [and others]. 1994. Dogwood anthracnose and its spread in the South. Prot. Rep. R8-PR-26. Atlanta, GA: U.S. Department of Agriculture, Forest Service, Southern Region. 10 p.
- Anderson, R.M. 1940. The spread of cottontail rabbits in Canada. *Canadian Field-Naturalist*. 54:70–72.
- Annand, E.M.; Thompson, F.R., III. 1977. Forest bird response to regeneration practices in central hardwood forests. *Journal of Wildlife Management*. 61: 159–171.
- Appel, David N.; Billings, Ronald F. 1992. Oak wilt perspectives. In: Proceedings of the national oak wilt symposium; 1992 June 22–25; Austin, TX. College Station, TX: Texas Agricultural Experiment Station, Texas Forest Service, Texas Agricultural Extension Service. 217 p.
- Aust, W.M.; Hodges, J.D.; Johnson, R.L. 1985. The origin, growth and development of natural, pure, even-aged stands of bottomland oak. In: Shoulders E., ed. Proceedings of the third biennial southern silvicultural research conference; 1984 November 7–8; Atlanta, GA. Gen. Tech. Rep. SO-54. New Orleans, LA: U. S. Department of Agriculture, Forest Service, Southern Forest Experiment Station: 163–170.
- Averill, Robert D.; Larson, Louise; Saveland, James M. [and others]. 1994. Disturbance processes and ecosystem management. [Available on the Internet at <<http://svinet2.fs.fed.us/research/vmpr/disturb.htm>>.] Washington, DC: U.S. Department of Agriculture, Forest Service. 33p.
- Bailey, Robert G; Avers, Peter E.; King, Thomas; McNab, W. Henry, eds. 1994. Ecoregions and subregions of the United States (map). Washington, DC. U.S. Department of the Interior, Geological Survey. Scale 1:7,500,000.
- Baker, J.B. 1992. Natural regeneration of shortleaf pine. In: Brissette, J.C.; Barnett, J.P., comps. Proceedings of the shortleaf pine regeneration workshop; 1991 October 29–31; Little Rock, AR. Gen. Tech. Rep. SO-90. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station: 102–112.
- Baker, J.B., tech. comp. 1994. Proceedings of the symposium on ecosystem management research in the Ouachita Mountains: pretreatment conditions and preliminary findings; 1993 October 26–27; Hot Springs, AR. Gen. Tech. Rep. SO-112. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 265 p.
- Baker, J.B.; Cain, M.D.; Guldin, J.M. [and others]. 1996. Uneven-aged silviculture for the loblolly and shortleaf pine forest cover types. Gen. Tech. Rep. SO-118. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Research Station. 65 p.
- Baldwin, F.L.; Boyd, J.W. 1998. Recommended chemicals for weed and brush control in Arkansas. Misc. Pub. 44. Little Rock, AR: University of Arkansas, Division of Agriculture, Cooperative Extension Service. 137 p.
- Baldwin, Ford L.; Boyd, John W.; Tripp, Timothy. 1988. Recommended chemicals for weed and brush control. Misc. Pub. 44. Little Rock, AR: University of Arkansas, Cooperative Extension Service; Fayetteville, AR: U.S. Department of Agriculture, Forest Service. 98 p.
- Barnett, J.P. 1992. Production of shortleaf pine seedlings. In: Brissette, J.C.; Barnett, J.P., comps. Proceedings of the shortleaf pine regeneration workshop; 1991 October 29–31; Little Rock, AR. Gen. Tech. Rep. SO-90. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station: 20–31.
- Barnett, J.P.; Brissette, J.C.; Carlson, W.C. 1986. Artificial regeneration of shortleaf pine. In: Murphy, P.A., ed. Proceedings of the symposium on the shortleaf pine ecosystem; 1986 March 31–April 2; Little Rock, AR. Monticello, AR: University of Arkansas, Department of Forest Resources, Cooperative Extension Service: 64–88.

- Barrett, J.W., ed. 1995. Regional silviculture of the United States. 3d ed. New York: John Wiley and Sons. 643 p.
- Bassett, E.N.; Fenn, P.; Mead, M.A. 1982. Drought-related mortality and incidence of *Hypoxylon* canker. Arkansas Farm Research. 31(1): 8.
- Batek, Michael J. 1994. Presettlement vegetation of the Current River watershed in the Missouri Ozarks. Columbia, MO: University of Missouri-Columbia. 264 p. M.S. thesis.
- Beck, D.E. 1988. Clearcutting and other regeneration options for upland hardwoods. In: Proceedings of the sixteenth annual hardwood symposium of the Hardwood Research Council: hardwood supply—feast and famine; 1988 May 15–18; Cashiers, NC. Cashiers, NC: Hardwood Research Council: 44–54.
- Beck, D.E.; Hooper, R.M. 1986. Development of a southern Appalachian hardwood stand after clearcutting. Southern Journal of Applied Forestry. 10: 168–172.
- Beilman, A.P.; Brenner, L.G. 1951. The recent intrusion of forests in the Ozarks. Annals of the Missouri Botanical Garden. 38: 261–282.
- Bey, Calvin F. 1964. Advance oak reproduction grows fast after clearcutting. Journal of Forestry. 62: 339–340.
- Bey, Calvin F. 1990. *Ulmus americana* L. American elm. In: Burns, Russell M.; Honkala, Barbara H., tech. coords. Silvics of North America, Volume 2, Hardwoods. Agric. Handb. 654. Washington, DC: U.S. Department of Agriculture, Forest Service: 801–807.
- Birdsey, Richard A.; May, Dennis M. 1988. Timber resources of east Oklahoma. Resour. Bull. SO-135. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 29 p.
- Boggs, J.A.; Wittwer, R.F. 1993. Emergence and establishment of shortleaf pine seeds under various seedbed conditions. Southern Journal of Applied Forestry. 17(1): 44–48.
- Bradbury, John. 1817. Travels in the interior of North America in the years 1809, 1810, 1811. London: Smith, Sherwood, Neely, and Jones. 346 p.
- Braun, E. Lucy. 1950. Deciduous forests of eastern North America. New York: Hafner. 596 p.
- Brawn, J.D.; Robinson, S.K. 1996. Source-sink population dynamics may complicate the interpretation of long-term census data. Ecology. 77: 3–12.
- Brennan, L.A. 1991. How can we reverse the northern bobwhite decline? Wildlife Society Bulletin. 19: 544–555.
- Brinkman, K.A.; Rogers, N.F. 1967. Timber management guide for shortleaf pine and oak-pine types in Missouri. Res. Pap. NC-19. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 15 p.
- Brissette, J.C.; Barnett, J.P., comps. 1992. Proceedings of the shortleaf pine regeneration workshop; 1991 October 29–31; Little Rock, AR. Gen. Tech. Rep. SO-90. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 236 p.
- Brissette, J.C.; Carlson, W.C. 1992. Seedling quality and field performance. In: Brissette, J.C.; Barnett, J.P., comps. Proceedings of the shortleaf pine regeneration workshop; 1991 October 29–31; Little Rock, AR. Gen. Tech. Rep. SO-90. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station: 32–43.
- Brown, Michael E. 1997. Developing a funding strategy for areas without existing gypsy moth populations. In: Proceedings of the 1997 annual gypsy moth review; 1997 Nov. 3–6; Charleston, WV. Charleston, WV: West Virginia Department of Agriculture: 114–118.
- Bukenhofer, George A.; Hedrick, Larry D. 1997. Shortleaf pine/bluestem grass ecosystem renewal in the Ouachita Mountains. Transactions of the 62nd North American Wildlife and Natural Resources Conference. 62: 509–515.
- Burns, Russell M.; Honkala, Barbara H. 1990. Silvics of North America, volume 1, conifers. Agric. Handb. 654. Washington, DC: U.S. Department of Agriculture, Forest Service. 498 p.
- Byther, R.S.; Davidson, R.M., Jr. 1979. Dogwood anthracnose. Ornamental Northwest Newsletter. 3: 20–21.
- Campbell, Faith T., ed. 1997. List of invasive exotic/alien/nonindigenous plant species [data base and printout]. Springfield, VA: National Coalition of Exotic Pest Councils. 25 p.
- Carter, M.F.; Barker, K. 1993. An interactive database for setting conservation priorities for western neotropical migrants. In: Finch, D.M.; Stangel, P.W., eds. Status and management of neotropical migratory birds. Gen. Tech. Rep. RM-229. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station: 120–127.
- Chandler, Craig ; Cheney, Philip; Thomas, Philip [and others]. 1983. Forest fire behavior and effects. In: Fire in forestry: Vol. I. New York: John Wiley and Sons. 450 p.
- Christisen, Donald M. 1970. Fox and gray squirrels. In: Nagel, W.O., ed. Conservation contrasts: three decades of

- non-political management of wildlife and forests in Missouri. Jefferson City, MO: Missouri Department of Conservation: 139–146.
- Clark, Alexander, III; Souter, R.A. 1996. Stem cubic-foot volume tables for tree species in the Arkansas area. Res. Pap. SE-297. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Research Station. 70 p.
- Clark, F.B. 1970. Measures necessary for natural regeneration of oaks, yellow-poplar, sweetgum, and black walnut. In: The silviculture of oaks and associated species. Res. Pap. NE-144. Upper Darby, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station: 1–16.
- Clark, F.B.; Boyce, S.G. 1964. Yellow-poplar seed remains viable in the forest litter. *Journal of Forestry*. 62(8): 564–567.
- Clark, Joseph D. 1988. Arkansas status report. In: Smith, Howard L.; de Almeida, Maria H., eds. Proceedings of the 9th eastern workshop on black bear research and management [Draft copy]. 1988 April 5–7; Huntsville, ON, Canada. Toronto, ON: Ontario Ministry of Natural Resources: 4–7. [Unpublished copy on file in the office of the Resources Specialist, U.S. Department of Agriculture, Forest Service, Mark Twain National Forest, Rolla, MO.]
- Cogbill, Charles V. 1995. Toward an ecological definition of eastern old growth. [Abstract]. In: Twenty-second annual natural areas conference, paper and poster abstracts; 1995 October 25–28; Fayetteville, AR. Little Rock, AR: Natural Areas Association; Arkansas Natural Heritage Commission; and The Nature Conservancy, Arkansas Field Office: 7.
- Colorado Bird Observatory. [Accessed November 1998]. Partners in Flight scores for physiographic region 19. <<http://members.aol.com/CBOPIFDB/index.html>>. [Various pagination.] (This Web site has since moved to <<http://www.cbobirds.org>>.)
- Conner, Michael D.; Wilkinson, Robert C. 1979. Ips bark beetles in the South. For. Insect & Dis. Leaf. 129. Washington, DC: U.S. Department of Agriculture, Forest Service. 8 p.
- Critchfield, W.B.; Little, E.L., Jr. 1966. Geographic distribution of the pines of the world. Misc. Pub. 991. Washington, DC: U.S. Department of Agriculture, Forest Service. 97 p.
- Croneis, C. 1930. Geology of the Arkansas Paleozoic era. Bull. 3. Little Rock, AR: Arkansas Geological Survey. 457 p.
- Cutter, B.E.; Guyette, R.P. 1994. Fire frequency on an oak-hickory ridgetop in the Missouri Ozarks. *American Midland Naturalist*. 132: 393–398.
- Dale, M.E.; Sonderman, D.L. 1984. Effect of thinning and potential quality of young white oak crop trees. Res. Pap. NE-539. Radnor, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station. 11 p.
- Daniel, T.W.; Helms, J.A.; Baker, F.S. 1979. Principles of silviculture. 2nd ed. New York: McGraw-Hill. 500 p.
- Daughtrey, M.L.; Hibben, C.R. 1983. Lower branch dieback, a new disease of northeastern dogwoods. *Phytopathology*. 73: 365.
- deCalesta, D.S.; Stout, S.L. 1997. Relative deer density and sustainability: a conceptual framework for integrating deer management with ecosystem management. *Wildlife Society Bulletin*. 25(2): 252–258.
- Delcourt, H.R.; Delcourt, P.A. 1991. Late-quaternary vegetation history of the Interior Highlands of Missouri, Arkansas, and Oklahoma. In: Henderson, Douglas; Hedrick, L.D., eds. Proceedings of the conference on the restoration of old growth forests in the Interior Highlands of Arkansas and Oklahoma; 1991 September 19–20; Morrilton, AR. Hot Springs, AR: U.S. Department of Agriculture, Forest Service, Ouachita National Forest; Morrilton, AR: Winrock Institute for Agricultural Development: 15–30.
- Della-Bianca, L.; Beck, D.E. 1985. Selection management in southern Appalachian hardwoods. *Southern Journal of Applied Forestry*. 9(3): 191–196.
- Derr, H.J.; Mann, W.F. 1971. Direct-seeding pines in the South. Agric. Handb. 391. Washington, DC: U.S. Department of Agriculture, Forest Service. 68 p.
- Dey, D.C.; Johnson, P.S.; Garrett, H.E. 1996. Modeling the regeneration of oak stands in the Missouri Ozark Highlands. *Canadian Journal of Forestry Research*. 26: 573–583.
- Dickson, J.G.; Thompson, F.R., III; Conner, R.N.; Franzreb, K.E. 1995. Silviculture in central and southeastern oak-pine forests. In: Martin, T.E.; Finch, D.M., eds. Ecology and management of neotropical migratory birds. New York: Oxford University Press: 245–268.
- Dixon, R.K.; Garrett, H.E.; Cox, G.S. 1979. Containerized shortleaf pine seedlings show superior growth and ectomycorrhizal development with mist foliar fertilization. *Southern Journal of Applied Forestry*. 3(4): 154–157.
- Doane, Charles C.; McManus, Michael L., eds. 1981. The gypsy moth: research toward integrated pest management. Tech. Bull. 1584. Washington, DC: U.S. Department of Agriculture, Forest Service. 757 p.
- Donovan, T.M.; Thompson, F.R., III; Faaborg, J.; Probst, J. 1995. Reproductive success of migratory birds in habitat

- sources and sinks. *Conservation Biology*. 9: 1380–1395.
- Dorman, K.W. 1976. The genetics and breeding of southern pines. *Agric. Handb.* 471. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 407 p.
- Drummond, David B. 1990. [Entomologist, U.S. Department of Agriculture, Forest Service, Southern Region, Forest Health Protection, Pineville, LA.] Personal communication. Hot Springs, AR: Ouachita National Forest. [Not paged]. [Unpublished report on file with the Entomologist, U.S. Department of Agriculture, Forest Service, Southern Region, Forest Health Protection. Pineville, LA.]
- Drummond, David B. 1992. [Entomologist, U.S. Department of Agriculture, Forest Service, Southern Region, Forest Health Protection, Pineville, LA.] Personal communication. Hot Springs, AR: Ozark National Forest. [Not paged]. [Unpublished report on file with the Entomologist, U.S. Department of Agriculture, Forest Service, Southern Region, Forest Health Protection. Pineville, LA.]
- du Pratz, L.P. 1774. *The History of Louisiana*. London: T. Becket. 405 p.
- Durham, J.F.; Kurtz, W.B.; Garrett, H.E. 1983. The silvicultural and economic feasibility of thinning mixed stands of oak on a site index of 70 in south Missouri. In: Muller, R.N., ed. *Proceedings, fourth central hardwood forest conference; 1982 November 8–10; Lexington, KY*. Lexington, KY: University of Kentucky: 25–31.
- Durner, G.M.; Gates, J.E. 1993. Spatial ecology of black rat snakes on Remington Farms, Maryland. *Journal of Wildlife Management*. 57: 812–826.
- Dwyer, John P.; Kurtz, William B. 1991. The realities of sustainable management vs. diameter limit harvest. *Northern Journal of Applied Forestry*. 8(4): 174–176.
- Earles, J.M. 1976. Forest statistics for east Oklahoma counties. *Resour. Bull.* SO-62. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 40 p.
- Ellison, Lincoln. 1960. Influence of grazing on plant succession of rangelands. *Botanical Review*. 26: 1–78.
- Essex, Burton L.; Spencer, John S., Jr. 1974. Timber Resources of Missouri's eastern Ozarks. *Resour. Bull.* NC-19. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 62 p.
- Farrar, Robert M. 1981. Regulation of uneven-aged loblolly-shortleaf pine forests. In: Barnett, J.P., ed. *Proceedings of the first biennial southern silvicultural research conference; 1980 November 6–7; Atlanta, GA*. Gen. Tech. Rep. SO-34. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station: 294–304 p.
- Featherstonhaugh, G.W. 1835. Geological report of an examination made in 1834 of the elevated country between the Missouri and Red Rivers. Ex. Doc. No. 151. Washington, DC: U.S. Congress. 97 p.
- Featherstonhaugh, G.W. 1844. *Excursion through the slave States: from Washington on the Potomac, to the frontier of Mexico; with sketches of popular manners and geological notices*. New York: Harper and Bros. 2 vol.
- Fenneman, Nevin M. 1938. *Physiography of Eastern United States*. New York: McGraw-Hill. 714 p.
- Ferguson, E.R.; Graney, D.L. 1972. Site index table for shortleaf pine in the Boston Mountains of Arkansas. *Res. Note* SO-137. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 4 p.
- Ferguson, E.R.; Graney, D.L. 1975. Site index tables for shortleaf pine in the Ozark Highlands of northern Arkansas and southern Missouri. *Res. Note* SO-207. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 5 p.
- Finch, D.M. 1991. Population ecology, habitat requirements, and conservation of neotropical birds. Gen. Tech. Rep. RM-205. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station. 26 p.
- Finch, D.M.; Stangel, P.W. 1993. Status and management of neotropical migratory birds. Gen. Tech. Rep. RM-229. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station. 422 p.
- Fischer, Burnell C. 1981. Designing forest openings for the group selection method. In: Barnett, J.P., ed. *Proceedings of the first biennial southern silvicultural research conference; 1980 November 6–7; Atlanta, GA*. Gen. Tech. Rep. SO-34. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station: 274–277.
- Fitzgibbon, Bobbye. 1997. Arkansas gypsy moth project. *Forest Health Protection Update*. 1(3): 5–6.
- Fletcher, P.W.; McDermott, R.E. 1957. Influence of geologic parent material and climate on distribution of shortleaf pine in Missouri. *Research Bull.* 625. Columbia, MO: University of Missouri, College of Agriculture, Agricultural Experiment Station. 43 p.

- Ford-Robertson, F.C., ed. 1971. Terminology of forest science, technology practice and products. English language version. Washington, DC: Society of American Foresters. 349 p.
- Foti, Thomas L. 1974. Natural divisions of Arkansas. In: Arkansas Natural Area Plan. Little Rock, AR: Arkansas Department of Planning: 11–37.
- Foti, Thomas L. 1976. Arkansas, its land and people. Little Rock, AR: Arkansas Department of Education, Environmental and Conservation Office. 71 p.
- Foti, Thomas L.; Glenn, Susan M. 1991. The Ouachita Mountain landscape at the time of settlement. In: Henderson, Douglas; Hedrick, L.D., eds. Proceedings of the conference on restoration of old growth forests in the Interior Highlands of Arkansas and Oklahoma; 1991 September 19–20; Morrilton, AR. Hot Springs, AR: U.S. Department of Agriculture, Forest Service, Ouachita National Forest; Morrilton, AR: Winrock Institute for Agricultural Development: 49–65.
- Foti, T.; Blaney, M.; Li, X.; Smith, K.G. 1994. A classification system for the natural vegetation of Arkansas. Proceedings of the Arkansas Academy of Science. 48: 50–62.
- Fowells, H.A., comp. 1965. Silvics of forest trees of the United States. Agric. Handb. 271. Washington, DC: U.S. Department of Agriculture, Forest Service. 762 p.
- Franco, Peter A.; Miller, Patrick E.; Hartsell, Andrew J. 1993a. Forest statistics for northeast Oklahoma counties—1993. Resour. Bull. SO-174. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 30 p.
- Franco, Peter A.; Miller, Patrick E.; Hartsell, Andrew J. 1993b. Forest statistics for southeast Oklahoma counties—1993. Resour. Bull. SO-176. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 31 p.
- Freemark, K. E.; Dunning, J.B.; Hejl, S.J.; Probst, J.R. 1995. A landscape ecology perspective for research, conservation, and management. In: Martin, T.E.; Finch, D.M., eds. Ecology and management of neotropical migratory birds. New York: Oxford University Press: 381–427.
- Gaines, Glen; Arndt, Paul; Croy, Steve [and others]. 1997. Guidance for conserving and restoring old-growth forest communities on national forests in the southern region; report of the Region 8 Old-Growth Team. For. Rep. R8-FR 62. Atlanta, GA: U.S. Department of Agriculture, Forest Service, Southern Region. 122 p.
- Gansner, David A. 1966a. Timber resources of Missouri's northwestern Ozarks. Bull. B847. Columbia, MO: University of Missouri, Agricultural Experiment Station, and U. S. Department of Agriculture, Forest Service. 25 p.
- Gansner, David A. 1966b. Timber resources of Missouri's southwestern Ozarks. Bull. B845. Columbia, MO: University of Missouri, Agricultural Experiment Station, and U. S. Department of Agriculture, Forest Service. 21 p.
- Gardner, J.E.; Gardner, T.L. 1982. An inventory and evaluation of cave resources of Mark Twain National Forest. [Final report submitted to the Mark Twain National Forest, USDA, in compliance with cooperative cave inventory agreement.] Rolla, MO: Missouri Department of Conservation. 100 p.
- Gerstacker, Friedrich. 1881. Wild sports in the far West. New York: J.W. Lovell. 409 p.
- Gilfillan, Corinna. 1994. Exotic pests: a growing threat to the environment. Washington, DC: National Audubon Society. 22 p.
- Gingrich, S.F. 1967. Measuring and evaluating stocking and stand density in upland hardwood forests in the Central States. Forest Science. 13: 38–53.
- Gingrich, S.F. 1970. Effects of density, thinning, and species composition on the growth and yield of eastern hardwoods. In: The silviculture of oaks and associated species. Res. Pap. NE-144. Upper Darby, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station: 26–35.
- Gottschalk, K.W. 1983. Management strategies for successful regeneration: oak-hickory. In: Proceedings of 1983 Penn State forestry issues conference—regenerating hardwood stands; 1983 March 15–16; University Park, PA. University Park, PA: Penn State University, School of Natural Resources: 190–213.
- Gottschalk, K.W. 1985. Effects of shading on growth and development of northern red oak, black oak, black cherry, and red maple seedlings: I. Height, diameter, and root/shoot ratio. In: Dawson, J.O.; Majerus, K.A., eds. Proceedings of the fifth central hardwood forest conference; 1985 April 15–17; Urbana, IL. Urbana-Champaign, IL: University of Illinois, Department of Forestry: 189–195.
- Gottschalk, Kurt W. 1990. Gypsy moth effects on mast protection. In: McGee, Charles E., ed. Proceedings, southern Appalachian mast management workshop; 1989 August 14–16; Knoxville, TN. Knoxville, TN: University of Tennessee: 42–50.

- Graney, D.L. 1976. Site index relationships for shortleaf pine and upland oaks in the Ozark-Ouachita Highlands of Missouri, Arkansas and Oklahoma. In: Fralish, J.S.; Weaver, G.T.; Schlesinger, R.C., eds. Central hardwood forest conference proceedings; 1976 October 17–19; Carbondale, IL. Carbondale, IL: Southern Illinois University: 309–326.
- Graney, D.L. 1989. Growth of oak, ash, and cherry reproduction following overstory thinning and understory control in upland hardwood stands of northern Arkansas. In: Miller, J.H., ed. Proceedings of the fifth biennial southern silvicultural research conference; 1988 November 1–3; Memphis, TN. Gen. Tech. Rep. SO-74. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station: 245–252.
- Graney, D.L.; Burkhart, H.E. 1973. Polymorphic site index curves for shortleaf pine in the Ouachita Mountains. Res. Pap. SO-85. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 14 p.
- Graney, D.L.; Ferguson, E.R. 1972. Shortleaf pine site-index relationships in the Ozark Highlands. Proceedings of the Soil Science Society of America. 36(3): 495–500.
- Graney, D.L.; Murphy, P.A. 1997. An evaluation of uneven-aged cutting methods in even-aged oak-hickory stands in the Boston Mountains of Arkansas. In: Pallardy, Stephen G.; Cecich, Robert A.; Garrett, H. Eugene; Johnson, Paul S., eds. Proceedings, 11th central hardwood forest conference; 1997 March 23–26; Columbia, MO. Gen. Tech. Rep. NC-188. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station: 130–146.
- Graney, D.L.; Rogerson, T.L. 1985. Development of oak, ash and cherry reproduction following thinning and fertilization of upland hardwood stands in the Boston Mountains of Arkansas. In: Shoulders, E., ed. Proceedings of the third biennial southern silvicultural research conference; 1984 November 7–8; Atlanta, GA. Gen. Tech. Rep. SO-54. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station: 171–177.
- Grazing Lands Forum. 1993. An explosion in slow motion: noxious weeds and invasive alien plants on grazing lands. Washington, DC: Grazing Lands Forum. 69 p.
- Guldin, J. M.; Baker, J.B. 1998. Uneven-aged silviculture, southern style. *Journal of Forestry*. 96: 22–26.
- Guldin, J.M.; Baker, J.B.; Shelton, M.G. 1994. Pretreatment conditions in mature pine/hardwood stands on south-facing slopes of the Ouachita/Ozark National Forests: midstory and overstory plants. In: Baker, J.B., tech. comp. Proceedings of the symposium on ecosystem management research in the Ouachita Mountains: pretreatment conditions and preliminary findings; 1993 October 26–27; Hot Springs, AR. Gen. Tech. Rep. SO-112. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 264 p.
- Guldin, J.M.; Carrillo, F.; Acosta, M.; Guries, R.P. 1998. Marking to cut versus marking to leave—a comparison of silvicultural tactics. In: Waldrop, T., tech. comp. Proceedings of the ninth biennial southern silvicultural research conference; 1997 February 25–27; Clemson, SC. Gen. Tech. Rep. SRS-20. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Research Station. 628 p.
- Guldin, J.M.; Shelton, M.G.; Wittwer, R.F. [and others]. 1993. New perspectives/ecosystem management research on the Ouachita/Ozark National Forests: phase II—silviculture research. Study Plan FS-SO-4106-91. Monticello, AR: U.S. Department of Agriculture, Forest Service, Southern Research Station. 64 p.
- Guyette, Richard P.; Cutter, Bruce E. 1991. Tree ring analysis of fire history in a post oak savanna in the Missouri Ozarks. *Natural Areas Journal*. 11: 93–99.
- Guyette, Richard P.; Cutter, Bruce E. 1997. Fire history, population, and calcium cycling in the Current River watershed. In: Pallardy, Stephen G.; Cecich, Robert A.; Garrett, H. Eugene; Johnson, Paul S., eds. Proceedings, 11th central hardwood forest conference; 1997 March 23–26; Columbia, MO. Gen. Tech. Rep. NC-188. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station: 355–372.
- Guyette, Richard P.; McGinnes, E.A. 1982. Fire history of an Ozark glade in Missouri. *Transactions of the Missouri Academy of Science*. 16: 85–93.
- Hagan, J.M., III; Johnston, D.W. 1992. Ecology and conservation of neotropical migrant landbirds. Washington, DC: Smithsonian Institution Press. 609 p.
- Haley, T.J. 1996. [Entomologist, U.S. Department of Agriculture, Forest Service, Southern Research Station, Pineville, LA.] Forest health evaluation of the Ouachita National Forest. Pineville, LA: U.S. Department of Agriculture, Forest Service, State and Private Forestry, Forest Health Protection: 23 p. [Unpublished report on file with the Entomologist, U.S. Department of Agriculture, Forest Service, Southern Research Station, Pineville, LA.]
- Hallgren, S.W.; Ferris, D.M. 1995. Benomyl applied to roots improves second-year survival and growth of shortleaf pine. *Southern Journal of Applied Forestry*. 19(1): 36–41.

- Hallgren, S.W.; Tauer, C.G. 1989. Root growth potential, first-year survival, and growth of shortleaf pine seedlings show effects of lift date, storage, and family. *Southern Journal of Applied Forestry*. 13(4): 163–169.
- Hallgren, S.W.; Tauer, C.G.; Weeks, D.L. 1993. Cultural, environmental, and genetic factors interact to affect performance of planted shortleaf pine. *Forest Science*. 39(3): 478–498.
- Halliday, William R. 1982. American caves and caving, techniques, pleasures, and safeguards of modern cave exploration. New York: Barnes and Noble. 348 p.
- Halls, Lowell K. 1977a. Flowering dogwood/*Cornus florida* L. In: Halls, Lowell K., ed. Southern fruit-producing woody plants used by wildlife. Gen. Tech. Rep. SO-16. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station: 33–35.
- Halls, Lowell K. 1977b. Chinkapin/*Castanea* spp. In: Halls, Lowell, K., ed. Southern fruit-producing woody plants used by wildlife. Gen. Tech. Rep. SO-16. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station: 144–145.
- Halls, Lowell K. 1984. White-tailed deer: ecology and management. Harrisburg, PA: Stackpole Books: 870 p.
- Hamel, P.B. 1992. The land manager's guide to the birds of the South. Chapel Hill, NC: The Nature Conservancy. 437 p.
- Hansen, Mark H.; Frieswyk, Thomas; Glover, Joseph F.; Kelly, John F. 1992. The eastwide forest inventory data base: user's manual. Gen. Tech. Rep. NC-151. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 48 p.
- Harlow, William M.; Harrar, Ellwood S. 1969. Textbook of dendrology. 5th ed. New York: McGraw-Hill: 285–287.
- Harmon, Debbie; Lockhart, Jami J.; Sabo, George, III. 1996. From surveyor's chains to satellites: a study of historic environmental change and land use on the Lee Creek Unit of the Ozark National Forest, Crawford and Washington Counties, Arkansas. Arkansas Archeological Survey Project 955. Fayetteville, AR: Arkansas Archeological Survey. 56 p.
- Harrington, C.A.; Brissette, J.C.; Carlson, W.C. 1989. Root system structure in planted and seeded loblolly and shortleaf pine. *Forest Science*. 35(2): 469–480.
- Harty, Francis M. 1991. How Illinois kicked the exotic habit. In: McKnight, Bill, N., ed. Proceedings of a conference on biological pollution: the control and impact of invasive exotic species; 1991 October 25–26; Indianapolis, IN. Indiana Academy of Science, Indianapolis, IN. 195–209.
- Harvey, Michael J.; Redman, Ronald K. 1996. Endangered bats of Arkansas: distribution, status, and ecology (1995–1996), annual report to Arkansas Game and Fish Commission. Proj. W-56-R. Little Rock, AR: Arkansas Game and Fish Commission. 29 p.
- Hatcher, Robert D., Jr.; Thomas, William A.; Viele, George W., eds. 1990. The Appalachian-Ouachita orogen in the United States. Boulder, CO: Geologic Society of America. 781 p.
- Hawker, Jon L. [n.d.] Missouri landscapes: a tour through time. Rolla, MO: Missouri Department of Natural Resources, Division of Geology and Land Survey. 326 p.
- Hayes, Jane L.; Strom, B.L.; Roton, L.M.; Ingram, L.L., Jr., inventors; U.S. Department of Agriculture, Forest Service and Mississippi State University, assignees. 1995. A repellent of bark beetles. U.S. Patent 5,403,863.
- Hazell, Don B. 1967. Effects of grazing intensity on plant composition, vigor, and production. *Journal of Range Management*. 20(4): 249–252.
- Heath, Darrel R.; Saugey, D.A.; Heidt, G.A. 1986. Abandoned mine fauna of the Ouachita mountains, Arkansas: vertebrate taxa. *Proceedings Arkansas Academy of Science*. 40: 33–36.
- Hedlund, Arnold; Earles, J.M. 1970. Forest statistics for Arkansas counties. Resour. Bull. SO-22. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 52 p.
- Henderson, Douglas; Hedrick, Larry D., eds. 1991. Proceedings of the conference on restoration of old growth forests in the Interior Highlands of Arkansas and Oklahoma; 1990 September 19–20; Morrilton, AR. Hot Springs, AR: U.S. Department of Agriculture, Forest Service, Ouachita National Forest; Morrilton, AR: Winrock International Institute for Agricultural Development. 190 p.
- Hibben, C.R.; Daughtrey, M.L. 1988. Dogwood anthracnose in Northeastern United States. *Plant Disease*. 72: 19–203.
- Hines, F. Dee. 1988a. Forest statistics for Arkansas' Ouachita counties—1988. Resour. Bull. SO-137. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 28 p.
- Hines, F. Dee. 1988b. Forest statistics for Arkansas' Ozark counties—1988. Resour. Bull. SO-131. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 39 p.
- Hoffman, Randy; Kearns, Kelly. 1997. Wisconsin manual of control recommendations for ecologically invasive plants. Madison, WI: Wisconsin Department of Natural Resources. 102 p.

- Horst, R. Kenneth. 1990. Westcott's plant disease handbook. 5th ed. New York: Chapman & Hall. 175.
- Houck, Louis. 1908. History of Missouri from the earliest explorations and settlements until the admission of the State into the union. Chicago: R.R. Donnelley & Sons. 3 Vols.
- Hu, S.; Burns, P.Y. 1987. Shortleaf pine: a bibliography, 1896–1984. Res. Rep. No. 11, April 1987. Baton Rouge, LA: Louisiana State University, Louisiana Agricultural Experiment Station, School of Forestry, Wildlife, and Fisheries. 97 p.
- Hubbell, Sue. 1993. Broadside from the other orders: a book of bugs. New York: Random House. 276 p.
- Huebschmann, Michael M.; Lynch, Thomas B.; Murphy, Paul.A. 1997. Shortleaf pine stand simulator: an even-aged natural shortleaf pine growth and yield model. Stillwater, OK: Oklahoma State University, Oklahoma Agricultural Experiment Station Research Report P-967. 25 p.
- Hunter, Malcolm L., Jr. 1996. Fundamentals of conservation biology. Cambridge, MA: Blackwell Science, Inc. 482 p.
- Hunter, W.C.; Carter, M.F.; Pashley, D.N.; Barker, K. 1993. The Partners in Flight species prioritization scheme. In: Finch, D. M.; Stangel, P. W., eds. Status and management of neotropical migratory birds. Gen. Tech. Rep. RM-229. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station: 109–119.
- Jacobson, Robert B. 1995. Spatial controls on patterns of landuse induced stream disturbance at the drainage-basin scale—an example from gravel-bed streams of the Ozark Plateaus, Missouri. *Geomorphology*. 89: 219–239.
- James, Douglas A.; Neal, Joseph C. 1986. Arkansas birds. Fayetteville, AR: The University of Arkansas Press. 402 p.
- James, E., ed. 1823. An expedition from Pittsburgh to the Rocky Mountains. Philadelphia: H.C. Carey. 442 p. + atlas. Vol. 2.
- Janzen, G.C.; Hodges, J.D. 1987. Development of advanced oak regeneration as influenced by removal of midstory and understory vegetation. In: Phillips, D.R., comp. Proceedings of the fourth biennial southern silvicultural research conference; 1986 November 4–6; Atlanta, GA. Gen. Tech. Rep. SE-42. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station: 455–461.
- Jenkins, Sean E.; Guyette, Richard P.; Rebertus, Alan J. 1997. Vegetation-site relationships and fire history of a savanna-glade-woodland mosaic in the Ozarks. In: Pallardy, Stephen G.; Cecich, Robert A.; Garrett, H. Eugene; Johnson, Paul S., eds. Proceedings, 11th central hardwood forest conference; 1997 March 23–26; Columbia, MO. Gen. Tech. Rep. NC-188. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station: 184–201.
- Jensen, Peter N.; Schumacher, C.M. 1969. Changes in prairie plant composition. *Journal of Range Management*. 22(1): 57–60.
- Johnson, F.L.; Schnell, G.D. 1985. Wildland fire history and the effects of fire on vegetative communities at Hot Springs National Park, Arkansas. Norman, OK: Oklahoma Biological Survey. 49 p.
- Johnson, George P. 1988. Revision of *Castanea* sect. *Balanocastanon* (Fagaceae). *Journal of the Arnold Arboretum*. 69: 25–49.
- Johnson, P.S. 1977. Predicting oak stump sprouting and sprout development in the Missouri Ozarks. Res. Pap. NC-149. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 11 p.
- Johnson, P.S. 1979. Shoot elongation of black oak and white oak sprouts. *Canadian Journal of Forestry Research*. 9: 489–494.
- Johnson, P.S. 1993. Perspectives on the ecology and silviculture of oak-dominated forests in the Central and Eastern States. Gen. Tech. Rep. NC-153. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 28 p.
- Johnson, P.S. 1997. The silviculture of upland central hardwoods: 25 years of change. In: Meyer, D.A., ed. Proceedings of the twenty-fifth annual hardwood symposium: 25 years of hardwood silviculture: a look back and a look ahead. Cashiers, NC: Hardwood Research Council: 17–44.
- Johnson, P.S.; Jacobs, R.D.; Martin, A.J.; Godel, E.D. 1989. Regenerating northern red oak: three successful case histories. *Northern Journal of Applied Forestry*. 6(4): 174–178.
- Johnson, W.T. 1956. The Asiatic oak weevil and other insects causing damage to chestnut foliage in Maryland. *Journal Economic Entomology*. 49(5): 717–718.
- Kessler, Kenneth J., Jr. 1992. Oak decline on public lands in the central forest region. Res. Note NC-362. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station: 1–4.
- Keys, James E. Jr.; Carpenter, Constance A.; Hooks, Susan L. [and others] 1995. Ecological units of the Eastern United States, first approximation. Atlanta, GA: U.S. Department of Agriculture, Forest Service, Southern Region. 82 p.

- Kimmins, J.P. 1987. Forest ecology. New York: MacMillan. 531 p.
- Kowal, R.J. 1960. Southern pine beetle. [Pest leaflet 49.] Washington, DC: U.S. Department of Agriculture, Forest Service. 7 p.
- Kreiter, Scott D. 1995. Dynamics and spatial pattern of a virgin old-growth hardwood-pine forest in the Ouachita Mountains, Oklahoma, from 1896 to 1994. Stillwater, OK: Oklahoma State University. 141 p. M.S. thesis.
- Kress, M.W.; Baker, R.; Ursic, S.J. 1990. Chemistry response of two forested watersheds to acid atmospheric deposition. *Water Resources Bulletin*. 26(5): 747–756.
- Kurtz, W.B.; Garrett, H.E.; Williams, R.A. 1981. Young stands of scarlet oak in Missouri can be thinned profitably. *Southern Journal of Applied Forestry*. 5(1): 12–16.
- Ladd, Douglas. 1991. Reexamination of the role of fire in Missouri oak woodlands. In: Proceedings, oak woods management workshop; October 21–22, 1988. Charleston, IL: Eastern Illinois University: 76–80.
- Ladd, Douglas; Huemann, Blane. 1994. [Director of Science, Stewardship and Registry, Missouri Field Office, The Nature Conservancy, St. Louis, MO.] Baseline ecological assessment of selected oak woodlands on the Houston-Rolla District, Mark Twain National Forest. Rolla, MO: U.S. Department of Agriculture, Forest Service. 66 p. [Unpublished report on file at the Supervisor's Office, U.S. Department of Agriculture, Forest Service, Mark Twain National Forest, Rolla, MO.]
- Lambeth, C.C. 1984. Large-scale planting of North Carolina loblolly pine in Arkansas and Oklahoma: a case of gain versus risk. *Journal of Forestry* 82(12): 736–741.
- Lanier, G.N. 1972. Biosystematics of the genus *Ips* (Coleoptera: Scolytidae) in North America, Happing's groups IV and X. *Canadian Journal of Entomology*. 49(5): 717–718.
- Larsen, D.R.; Metzger, M.A.; Johnson, P.S. 1997. Oak regeneration and overstory density in the Missouri Ozarks. *Canadian Journal of Forest Research*. 27(6): 869–75.
- LaVal, R.K.; LaVal, M.L. 1980. Ecological studies and management of Missouri bats, with emphasis on cave-dwelling species. Terrestrial ser. No. 8. Columbia, MO: Missouri Department of Conservation. 53 p.
- Law, J.R.; Gott, J.D. 1987. Oak mortality in the Missouri Ozarks. In: Hay, Ronald L.; Woods, Frank W.; DeSelm, Hal, eds. Proceedings, central hardwood forest conference VI; 1987 February 24–26; Knoxville, TN. Knoxville, TN: University of Tennessee, Division of Continuing Education: 427–436.
- Law, J.R.; Lorimer, C.G. 1989. Managing uneven-aged stands. In: Clark, F. Bryan; Hutchinson, Jay G., eds. Central hardwood notes. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station: 6.08-1–6.08-6.
- Lawson, E.R. 1990. *Pinus echinata* Mill, shortleaf pine. In: Burns, R.M.; Honkala, B.H.; tech. coords. *Silvics of North America: Volume 1, Conifers*. Agric. Handb. 654. Washington, DC: U.S. Department of Agriculture, Forest Service, Timber Management Research: 316–326.
- Lawson, E.R.; Kitchens, R. 1983. Shortleaf pine. In: Burns, R.M., tech. comp. *Silvicultural systems for the major forest types of the United States*. Agric. Handb. 445. Washington, DC: U.S. Department of Agriculture, Forest Service: 157–161.
- Laycock, William A. 1967. How heavy grazing and protection affect sagebrush-grass ranges. *Journal of Range Management*. 20(4): 206–213.
- Leatherberry, Earl C. 1990. Timber resources of Missouri's eastern Ozarks. Resour. Bull. NC-115. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 48 p.
- Lee, J. Charles. 1980. Integrated use in the South: a biological perspective. In: Child, R. Dennis; Byington, Everet, eds. Proceedings, southern forest range and pasture symposium; 1980 March 13–14; New Orleans, LA. Morrilton, AR: Winrock International: 51–57.
- Leopold, Aldo. 1931. Report on a game survey of North Central States. Madison, WI: Democrat Printing: 299 p.
- Lewis, J.B. 1992. Eastern wild turkey in Midwestern oak-hickory forest. In: Dickson, James G., ed. *The wild turkey: biology and management*. Harrisburg, PA: Stackpole Books: 286–305.
- Liebhold, A.M.; Halverson, J.A.; Elmes, G.A. 1992. Gypsy moth invasion in North America: a quantitative analysis. *Journal of Biogeography*. 19: 513–520.
- Little, Elbert L., Jr. 1977. Atlas of United States trees: minor eastern hardwoods. Misc. Pub. No. 1342. Washington, DC: U.S. Department of Agriculture, Forest Service. 17 p. + 230 maps. Vol. 4.
- Lockhart, Jami J.; Hilliard, Jerry E.; Sabo, George, III; Weddle, Deborah, A. 1995. The evolution of human ecosystems in the Ozark National Forest, a pilot study of the Lee Creek Unit. Arkansas Archeological Survey Project 876. Fayetteville, AR: Arkansas Archeological Survey. 125 p.
- Loewenstein, E.F. 1996. An analysis of the size- and age-structure of an uneven-aged oak forest. Columbia, MO: University of Missouri. 167 p. Ph.D. dissertation.

- Loewenstein, E.F.; Garrett, H.E.; Johnson, P.S.; Dwyer, J.P. 1995. Changes in a Missouri Ozark oak-hickory forest during 40 years of uneven-aged management. In: Gottschalk, K.W.; Fosbroke, S.L.C., eds. Proceedings, 10th central hardwood forest conference; 1995 March 5–8; Morgantown, WV. Gen. Tech. Rep. NE-197. Radnor, PA: U. S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station: 159–164.
- Loftis, D.L. 1983. Regenerating red oak on productive sites in the southern Appalachians: a research approach. In: Jones, E.P., ed. Proceedings of the second biennial southern silvicultural research conference; 1982 November 4–5; Atlanta, GA. Gen. Tech. Rep. SE-24. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station: 144–150.
- Loftis, D.L. 1990. A shelterwood method for regenerating red oak in the southern Appalachians. *Forest Science*. 36(4): 917–929.
- London, Jack D. 1997. Forest statistics for Arkansas counties—1995. Resour. Bull. SRS-17. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 86 p.
- Lorimer, C.G. 1989. The oak regeneration problem: new evidence on causes and possible solutions. In: Proceedings, seventeenth annual hardwood symposium of the Hardwood Research Council; 1989 May 7–10; Merrimac, WI. Cashiers, NC: Hardwood Research Council: 23–40.
- Lorimer, C.G. 1992. Causes of the oak regeneration problem. In: Oak regeneration: serious problems and practical recommendations. Gen. Tech. Rept. SE-84. Asheville, NC: U.S. Department of Agriculture, Forest Service: 14–39.
- Lynch, T.B.; Murphy, P.A. 1995. A compatible height prediction and projection system for individual trees in natural, even-aged shortleaf pine stands. *Forest Science* 41(1): 194–209.
- Lynch, T.B.; Murphy, P.A.; Lawson, E.R. 1991. Stand volume equations for managed natural even-aged shortleaf pine [*Pinus echinata*] in eastern Oklahoma and western Arkansas. Res. Rept. P-921. Stillwater, OK: Oklahoma State University, Division of Agriculture, Oklahoma Agricultural Experiment Station. 12 p.
- MacDonald, W.L.; Hindal, D.F. 1981. Life cycle and epidemiology of *Ceratocystis*. In: Mace, Marshall E.; Bell, Alois A.; Beckman, Carl H., eds. Fungal wilt diseases of plants. New York: Academic Press: 128–139.
- Manion, P.D. 1981. Tree disease concepts. Englewood Cliffs, NJ: Prentice-Hall, Inc: 324–339.
- Marini, M.A.; Robinson, S.K.; Heske, E.J. 1995. Edge effects on nest predation in the Shawnee National Forest, southern Illinois. *Biological Conservation*. 74: 203–213.
- Marquis, D.A. 1965. Controlling light in small clearcuttings. Res. Pap. NE-39. Upper Darby, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station. 16 p.
- Marquis, D.A. 1978. Application of uneven-aged silviculture on public and private lands. In: U.S. Department of Agriculture. Uneven-aged silviculture and management in the United States: combined: Proceedings of two in-service workshops; 1975 July 15–17; Morgantown, WV; 1976 October 19–21; Redding, CA. Gen. Tech. Rep. WO-24. Washington, DC: U.S. Department of Agriculture, Forest Service: 25–63.
- Marquis, R.J.; Whelan, C.J. 1994. Insectivorous birds increase growth of white oak through consumption of leaf chewing insects. *Ecology*. 75: 2007–2014.
- Martin, B.; Kline, D. 1985. Fire management considerations and fire regimes. Rolla, MO: U.S. Department of Agriculture, Forest Service, Mark Twain National Forest. 12 p.
- Martin, T.E.; Finch, D.M. 1995. Ecology and management of neotropical migratory birds. New York: Oxford University Press. 489 p.
- Martin, Paul; Houf, Garry F. 1993. Glade grasslands in southwest Missouri. *Rangelands*. 15(2): 70–72.
- Masters, R.E.; Engle, D.M.; Robinson, R. 1993. Effects of timber harvest and periodic fire on soil chemical properties in the Ouachita Mountains. *Southern Journal of Applied Forestry*. 17(3): 139–145.
- Masters, R.E.; Skeen, John E.; Whitehead, James. 1995. Preliminary fire history of McCurtain County Wilderness Area and implications for red-cockaded woodpecker management. In: Kulhavy, David L.; Hopper, Robert G.; Costa, Ralph, eds. Red-cockaded woodpecker: species recovery, ecology, and management. Nacogdoches, TX: Stephen F. Austin University, Center for Applied Studies: 290–302.
- Masters, R.E.; Wilson, C.W.; Bukenhofer, G.A.; Payton, M.E. 1996. Effects of pine-grassland restoration for red-cockaded woodpeckers on white-tailed deer forage production. *Wildlife Society Bulletin*. 24(1): 77–84.
- Mattoon, W.B. 1915. Life history of shortleaf pine. Bull. 244. Washington, DC: U.S. Department of Agriculture, Forest Service. 46 p.
- McAndrews, A.H. 1926. The biology of the southern pine beetle. New York State College of Forestry, Syracuse, NY. 103 p. M.S. Thesis.

- McGarigal, Kevin; Marks, Barbara J. 1995. FRAGSTATS: spatial analysis program for quantifying landscape structure. Gen. Tech. Rep. PNW-GTR 351. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 122 p.
- McGee, C.E. 1984. Heavy mortality and succession in a virgin mixed mesophytic forest. Res. Pap. SO-209. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 9 p.
- McGinnes, E.A., Jr. 1982. Fire history of an Ozark glade in Missouri. Transactions Missouri Academy of Science. 16: 85–93.
- McKinley, Daniel. 1962. The history of the black bear in Missouri. Bluebird. 29(3): 1–15.
- McLemore, B.F. 1990. *Cornus florida* L. Flowering dogwood. In: Burns, Russell M.; Honkala, Barbara H., tech. coords. Silvics of North America: volume 2 hardwoods. Agric. Handb. 654. Washington, DC: U.S. Department of Agriculture, Forest Service: 278–283.
- McNab, W.H.; Avers, P.E., comps. 1994. Ecological subregions of the United States: section descriptions. WO-WSA-5. Washington, DC: U.S. Department of Agriculture, Forest Service. [Not paged].
- McQuilkin, R.A. 1975. Growth of four types of white oak reproduction after clearcutting in the Missouri Ozarks. Res. Pap. NC-116. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 5 p.
- Mendel, Joseph J. 1961. Timber resources of the eastern Ozarks. Bull. B779. Columbia, MO: University of Missouri, Agricultural Experiment Station; in cooperation with the U.S. Department of Agriculture, Forest Service. 27 p.
- Merz, R.W.; Boyce, S.G. 1956. Age of oak seedlings. Journal of Forestry. 54(11): 774–775.
- Meyer, John. 1986. Management of old growth forests in Missouri. Jefferson City, MO: State of Missouri, Conservation Commission. 16 p.
- Mielke, Manfred W.; Daughtrey, Margery L. 1991. How to identify and control dogwood anthracnose [Leaflet]. NA-GR-18. Radnor, PA: U.S. Department of Agriculture, Forest Service, Northeastern Area, State and Private Forestry. [Not paged].
- Miles, Patrick D. 1990. Timber resources of Missouri's southwest Ozarks. Resour. Bull. NC-116. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 44 p.
- Miller, G.W. 1996. Epicormic branching on central Appalachian hardwoods 10 years after deferment cutting. Res. Pap. NE-702. Radnor, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station. 9 p.
- Miller, James E.; Leopold, Bruce D. 1992. Population influences: predators. In: James G., ed. The wild turkey: biology and management. Harrisburg, PA: Stackpole Books: 119–128.
- Minckler, L.S.; Woerheide, J.D. 1965. Reproduction of hardwoods 10 years after cutting as affected by site and opening size. Journal of Forestry. 63(2): 103–107.
- Mitchell, R.J.; Garrett, H.E.; Cox, G.S. [and others]. 1987. Boron fertilization, ectomycorrhizal colonization, and growth of *Pinus echinata* seedlings. Canadian Journal of Forest Research. 17(10): 1153–1156.
- Monks, William. 1907. History of southern Missouri and northern Arkansas: being an account of the early settlements, the Civil War, the Ku-Klux, and times of peace. West Plains, MO: West Plains Journal Company. 247 p.
- Moser, John C.; Bridges, J. Robert; Yin, Hui-Fen. 1995. Ascospore dispersal of *Ceratocystiopsis ranaculosus*, a mycangial fungus of the southern pine beetle. Mycologia. 87(1): 84–86.
- Mudd, J.A. 1888. History of Lincoln County, Missouri from the earliest time to the present. Chicago, IL: Goodspeed. 261 p.
- Murphy, P.A. 1982. Sawtimber growth and yield for natural even-aged stands of shortleaf pine in the West Gulf. Res. Pap. SO-181. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 13 p.
- Murphy, P.A., ed. 1986. Proceedings of symposium on the shortleaf pine ecosystem; 1986 March 29–April 2; Little Rock, AR. Monticello, AR: University of Arkansas, Arkansas Cooperative Extension Service, Department of Forest Resources. 272 p.
- Murphy, P.A.; Beltz, R.C. 1981. Growth and yield of shortleaf pine in the West Gulf region. Res. Pap. SO-169. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 15 p.
- Murphy, P.A.; Baker, J.B.; Lawson, E.R. 1991. Selection management of shortleaf pine in the Ouachita Mountains. Southern Journal of Applied Forestry. 15(1): 61–67.
- Murphy, P.A.; Lawson, E.R.; Lynch, T.B. 1992. Basal area and volume development of natural even-aged shortleaf pine stands in the Ouachita Mountains. Southern Journal of Applied Forestry. 16(1): 30–34.
- Murphy, P.A.; Shelton, M.G.; Graney, D.L. 1993. Group selection—problems and possibilities for the more shade-tolerant

- species. In: Gillespie A.R.; Parker, G.R.; Pope, P.E.; Rink, G., eds. Proceedings, 9th central hardwood forest conference; Gen. Tech. Rep. NC-161. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station: 229–247.
- Mutch, Robert W.; Cook, Wayne A. 1996. Restoring fire to ecosystems: methods vary with land management goals. In: U.S. Department of Agriculture, Forest Service. The use of fire in forest restoration. Ogden, UT: U.S. Department of Agriculture, Forest Service, Intermountain Research Station: 9–11.
- Natural Areas Association. 1992. Compendium on exotic species. Mukwonago, WI: Natural Areas Association. 174 p.
- Nelson, John C. 1997. Presettlement vegetation patterns along the 5th Principal Meridian, Missouri Territory, 1815. *American Midland Naturalist*. 137: 79–94.
- Nelson, Paul. 1993. Quantum leap: back to a hidden paradise. *Missouri Resource Review*. Spring: 16–21.
- Nelson, Paul W. 1985. The terrestrial natural communities of Missouri. Jefferson City, MO: Missouri Department of Natural Resources. 197 p.
- Nickles, J.K.; Tauer, C.G.; Stritzke, J.F. 1981. Use of prescribed fire and hexazinone (Velpar) to thin understory shortleaf pine in an Oklahoma pine-hardwood stand. *Southern Journal of Applied Forestry*. 5(3): 124–127.
- Nigh, T.A.; Pallardy, S.G.; Garrett, H.E. 1985. Changes in upland oak-hickory forests of central Missouri: 1968–1982. In: Dawson, J.O.; Majerus, K.A., eds. Proceedings of the fifth central hardwood forest conference; 1985 April 15–17; Urbana, IL: University of Illinois, Department of Forestry: 170–177.
- Nigh, Timothy A.; Pflieger, William L.; Redfearn, Paul L. Jr. [and others]. 1992. The biodiversity of Missouri: definition, status, and recommendations for its conservation. [Report of the Biodiversity Task Force.] Jefferson City, MO: Missouri Department of Conservation and Mark Twain National Forest. 53 p.
- Noss, Reed F.; LaRoe, Edward T., III; Scott, J. Michael. 1995. Endangered ecosystems of the United States: a preliminary assessment of loss and degradation. Biological Rep. 28. Washington, DC: U.S. Department of the Interior, National Biological Service. 59 p.
- Novak, Milan; Baker, James; Obbard, Martyn; Malloch, Bruce, eds. 1987. Wild furbearer management and conservation in North America. Toronto: Ontario Ministry of Natural Resources. 1,150 p.
- Nowacki, Gregory J.; Trianowsky, Paul A. 1993. Literature on old-growth forests of Eastern North America. *Natural Areas Journal*. 13(2): 87–107.
- Nuttall, Thomas. 1837. Collections toward a flora of the territory of Arkansas. *Transactions of the American Philosophical Society*. 2: 139–202.
- Nuttall, Thomas. 1966. A journal of travels into the Arkansaw Territory. Ann Arbor, MI: University Microfilms. 296 p.
- Nuttall, Thomas. 1980. A journal of travels into the Arkansaw Territory during the year 1819. In: Lottinville, Savoie, ed. The American exploration and travel series. Norman, OK: University of Oklahoma Press. 361 p. Vol. 66.
- Nyland, R.D. 1996. Silviculture concepts and applications. New York: McGraw-Hill. 633 p.
- Oklahoma Biodiversity Task Force (OK BTF). 1996. Oklahoma's biodiversity plan: a shared vision for conserving our natural heritage. Oklahoma City, OK: Oklahoma Department of Wildlife Conservation. 129 p.
- Omernik, J.M. 1987. Ecoregions of the conterminous United States. *Annals of the Association of American Geographers*. 77(1): 118–125.
- Ostrom, Arnold J.; Hahn, Jerold T. 1974. Timber resources of Missouri's southwestern Ozarks. Resour. Bull. NC-21. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 60 p.
- Ostry, M.E.; Mielke, M.E.; Anderson, R.L. 1996. How to identify butternut canker and manage butternut trees. HT-70. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. [Not paged].
- Ostry, M.E.; Mielke, M.E.; Skilling, D.D. 1994. Butternut—strategies for managing a threatened tree. Gen. Tech. Rep. NC-165. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 7 p.
- Packard, S. 1991. Restoring oak ecosystems. *Restoration and Management Notes*. 11(1): 5–16.
- Pallardy, S.G.; Nigh, T.A.; Garrett, H.E. 1988. Changes in forest composition in central Missouri: 1968–1982. *American Midland Naturalist*. 120(2): 380–390.
- Palmer, Ernest J. 1921. The forest of the Ozark region. *Journal of the Arnold Arboretum*. 2: 216–232.
- Parker, G.R. 1989. Old-growth forests of the central hardwood region. *Natural Areas Journal*. 9(1): 5–11.
- Paton, P. 1994. The effect of edge on avian nest success: how strong is the evidence? *Conservation Biology*. 8: 17–26.

- Payne, T.L.; Billings, Ronald F. 1989. Evaluation of (S)-verbenone applications for suppressing southern pine beetle (Coleoptera: Scolytidae) infestations. *Journal of Economic Entomology*. 82(6): 1702–1708.
- Pell, William F. 1983. The natural divisions of Arkansas: a revised classification and description. *Natural Areas Journal*. 3(2): 12–23.
- Penfound, William T. 1964. The relation of grazing to plant succession in the tall grass prairie. *Journal of Range Management*. 17(5): 256–260.
- Pharris, L.D. 1981. Evaluation of black bear survey data in Arkansas, 1976–1980. *Proceedings of the Southeastern Association of Fish and Wildlife Agencies*. 35: 66–70.
- Probst, J.R.; Thompson, F.R., III. 1996. A multi-scale assessment of the geographic and ecological distribution of midwestern neotropical migratory birds. In: Thompson, F. R., III, ed. *Management of midwestern landscapes for the conservation of neotropical migratory birds*. Gen. Tech. Rep. NC-187. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station: 21–40.
- Puckette, W.L. 1974–75. Bear caves. *Oklahoma Underground*. 7: 15–17.
- Pyne, Stephen J. 1982. *Fire in America: a cultural history of wildland and rural fire*. Princeton, NJ: Princeton University Press. 654 p.
- Rafferty, M.D. 1996. Rude pursuits and rugged peaks, Schoolcraft's Ozark journal 1818–1819. Fayetteville, AR: The University of Arkansas Press. 170 p.
- Rebertus, A. 1994. A changing view of fire. *Missouri Wildlife*. Oct–Nov: 8.
- Renlund, D.W., ed. 1971. *Forest pest conditions in Wisconsin*. Ann. Rep. Madison, WI: Wisconsin Department of Natural Resources. 53 p.
- Rexrode, Charles O.; Brown, H. Daniel. 1983. Oak wilt. For. Insect & Dis. Leaf. 29. Washington, DC: U.S. Department of Agriculture, Forest Service. 6 p.
- Reynolds, R.R. 1969. Twenty-nine years of selection timber management on the Crossett Experimental Forest. Res. Pap. SO-40. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 19 p.
- Reynolds, R.R.; Baker, J.B.; Ku, T.T. 1984. Four decades of selection timber management on the Crossett Farm forestry forties. Bull. 872. Fayetteville, AR: University of Arkansas, Division of Agriculture, Arkansas Agricultural Experiment Station. 43 p.
- Rink, George. 1990. *Juglans cinerea* L. Butternut. In: Burns, Russell M.; Honkala, Barbara H., tech. coords. *Silvics of North America: volume 2, hardwoods*. Agric. Handb. 654. Washington, DC: U.S. Department of Agriculture, Forest Service: 386–390.
- Roach, B.A. 1962. Practical silviculture for central hardwood stands. *Southern Lumberman*: 34–38.
- Roach, B.A.; Gingrich, S.F. 1968. Even-aged silviculture for upland central hardwoods. Agric. Handb. 355. Washington, DC: U.S. Department of Agriculture, Forest Service. 39 p.
- Robbins, C. S.; Bystrak, D.; Geissler, P.H. 1986. The breeding bird survey: its first fifteen years, 1965–1979. Res. Pub. 157. Washington, DC: U.S. Fish and Wildlife Service. 196 p.
- Robinson, S.K.; Thompson, F.R., III; Donovan, T.M. [and others]. 1995. Regional forest fragmentation and the nesting success of migratory birds. *Science*. 26: 1987–1990.
- Rodgers, M.J. 1973. Movements and reproductive success of black bear introduced into Arkansas. *Proceedings of the annual conference of the Southeastern Association of Game and Fish Commissioners*. 27: 307–308.
- Roelofs, W. 1873. *Curculionides recueillis* au Japan par M.G. Lewis. *Annual Society Entomology de Belgique*. 16: 168.
- Rogers, J.G., Jr.; Dwyer, D.J.; Martin, C. 1993. Partners in flight: past, present, and future; a Government perspective. In: Finch, D.M.; Stangel, P.W., eds. *Status and management of neotropical migratory birds*. Gen. Tech. Rept. RM-229. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station: 5–6.
- Rosson, James F., Jr.; London, Jack D. 1997. Forest statistics for Arkansas' Ozark counties—1995. Resour. Bull. SRS-15. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 45 p.
- Rosson, James F., Jr.; London, Jack D. 1997. Forest statistics for Arkansas' Ouachita counties—1995. Resour. Bull. SRS-10. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Research Station. 39 p.
- Rothstein, S. I.; Verner, J.; Stevens, E. 1984. Radio-tracking confirms a unique diurnal pattern of spatial occurrence in the parasitic brown-headed cowbird. *Ecology*. 65: 77–88.
- Rowland, E.R., ed. 1930. *Life, letters and papers of William Dunbar*. Jackson, MS: Mississippi Historical Society. 410 p.
- Rudis, Victor A.; Tansey, John B. 1995. Regional assessment of remote forests and black bear habitat from forest resource surveys. *Journal of Wildlife Management*. 59(1): 170–180.

- Runkle, James R. 1991. Natural disturbance regimes and the maintenance of stable regional floras. In: Henderson, Douglas; Hedrick, Larry D., eds. Proceedings of the conference on the restoration of old growth forests in the Interior Highlands of Arkansas and Oklahoma; 1990 September 19–20; Morrilton, AR. Hot Springs, AR: U.S. Department of Agriculture, Forest Service, Ouachita National Forest; Morrilton, AR: Winrock International Institute for Agricultural Development: 31–48.
- Salwasser, H. 1994. Ecosystem management: can it sustain diversity and productivity? *Journal of Forestry*. 92: 6–10.
- Sander, I.L. 1966. Composition and distribution of hardwood reproduction after harvest cutting. In: Proceedings, 1966 symposium on hardwoods of the Piedmont and Coastal Plain. Macon, GA: Georgia Forest Research Council: 30–33.
- Sander, I.L. 1971. Height growth of new oak sprouts depends on size of advance reproduction. *Journal of Forestry*. 69(11): 809–811.
- Sander, I.L. 1977. Manager's handbook for oaks in the North Central States. Gen. Tech. Rep. NC-37. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 35 p.
- Sander, I.L.; Clark, F.B. 1971. Reproduction of upland hardwood forests in the Central States. *Agric. Handb.* 405. Washington, DC: U.S. Department of Agriculture, Forest Service. 25 p.
- Sander, I.L.; Graney, D.L. 1993. Regenerating oaks in the Central States. In: Loftis, D.L.; McGee, C.E., eds. Oak regeneration: serious problems, practical recommendations. Gen. Tech. Rep. SE-84. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station: 174–183.
- Sander, I.L.; Johnson, P.S.; Rogers, R. 1984. Evaluating oak advance reproduction in the Missouri Ozarks. Res. Pap. NC-251. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 16 p.
- Sander, I.L.; Johnson, P.S.; Watt, R.F. 1976. A guide for evaluating the adequacy of oak advance reproduction. Gen. Tech. Rep. NC-23. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 7 p.
- Sander, Ivan L. 1980. Some silvicultural and management options for upland hardwoods of the Mid-South. In: Proceedings, Mid-South upland hardwood symposium for the practicing forester and land manager; 1980 April 30–May 2; Harrison, AR. Tech. Pub. SA-TP 12. Atlanta, GA: U.S. Department of Agriculture, Forest Service, State and Private Forestry, Southeastern Area: 88–96.
- Sanders, Darryl. 1991. Bee and wasp pests. Columbia, MO: University of Missouri-Columbia, University Extension. 4 p.
- Sanderson, G.C. 1987. Raccoon. In: Novak, Milan; Baker, James A.; Obbard, Martyn E.; Malloch, Bruce, eds. Wild furbearer management and conservation in North America. Toronto, ON: Ontario Ministry of Natural Resources: 486–499.
- Sauer, J.R.; Hines, J.E.; Gough, G. [and others]. [Accessed January 1997]. The North American breeding bird survey: results and analysis. Version 96.3. <<http://www.mbr.nbs.gov>>. [Various pagination].
- Saughey, David A.; Heidt, Gary A.; Heath, Darrell R. 1988. Utilization of abandoned mine drifts and fracture caves by bats and salamanders: unique subterranean habitat in the Ouachita mountains. In: Proceedings of the symposium on management of amphibians, reptiles and small mammals in North America; 1998 July 19–21; Flagstaff, AZ. Gen. Tech. Rep. RM-166. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station: 64–71.
- Saughey, David A.; Heidt, Gary A.; Heath, Darrell R. 1989. Bats of the Ouachita Mountains. *Proceedings of the Arkansas Academy of Science*. 43: 33–36.
- Saveland, Jim. 1995. Fire in the forest. In: Eskew, Lane G., comp. Forest health through silviculture workshop; 1995 May 8–11; Mescalero, NM. Gen. Tech. Rep. RM-GTR-267. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station: 14–19.
- Schlesinger, R.C. 1978. Increased growth of released white oak poles continues through two decades. *Journal of Forestry*. 76(11): 726–727.
- Schlesinger, Richard C. 1976. Sixteen years of selection silviculture in upland hardwood stands. Res. Pap. NC-125. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 6 p.
- Schoolcraft, Henry Rowe. 1821. *Journal of a tour into the interior of Missouri and Arkansas performed in the year 1818 and 1819*. London: Richards Phillips and Co. 170 p.
- Schroeder, Walter A. 1982. Presettlement prairie of Missouri. *Natural History Series, No. 2*. Jefferson City: Missouri Department of Conservation. 37 p.
- Schroeder, Walter; Haithcoat, Tim; Porter, Shannon 1997. Pilot project proposal, Missouri historic vegetation project. [Unpublished report on file at the Forest Supervisor's office, U.S. Department of Agriculture, Forest Service, Mark Twain National Forest, Rolla, MO.]

- Schruben, P.G.; Arndt, R.E.; Bawiec, W.J.; Ambroziac, R.J. 1994. Geology of the conterminous U.S. at 1:2,500,000 scale [A digital representation of the P.B. King and H.M. Beikman 1974 map of geology of the conterminous United States]. U.S. Geological Survey, Oklahoma City, OK.
- Sealander, John A.; Heidt, Gary A. 1990. Arkansas mammals: their natural history, classification, and distribution. Fayetteville, AR: The University of Arkansas Press. 308 p.
- Shelton, M.G. 1997. Development of understory vegetation in pine and pine-hardwood shelterwood stands in the Ouachita Mountains—the first three years. Res. Pap. SRS-8. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Research Station. 18 p.
- Shelton, M.G.; Murphy, P.A. 1997. Understory vegetation 3 years after implementing uneven-aged silviculture in a shortleaf pine-oak stand. Res. Pap. SO-296. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Research Station. 13 p.
- Shelton, M.G.; Wittwer, R.F. 1992. Effects of seedbed condition on natural shortleaf pine regeneration. In: Brissette, J.C.; Barnett, J.P., comps. Proceedings of the shortleaf pine regeneration workshop; 1991 October 29–31; Little Rock, AR. Gen. Tech. Rep. SO-90. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station: 124–139.
- Shelton, M.G.; Wittwer, R.F. 1996. Shortleaf pine seed production in natural stands in the Ouachita and Ozark Mountains. Southern Journal of Applied Forestry. 20(2): 74–90.
- Shores, Elizabeth F. 1994. The red imported fire ant: mythology and public policy, 1957–1992. Arkansas Historical Quarterly. 53(3): 320–339.
- Sims, Daniel H. 1980. An overview of Mid-South upland hardwood regeneration problems and alternatives. In: Mid-South upland hardwood symposium for the practicing forester and land manager: Proceedings of a symposium; 1980 April 30–May 2; Harrison, AR. Tech. Pub. SA-TP 12. Atlanta, GA: U.S. Department of Agriculture, Forest Service, Division of State and Private Forestry, Southeastern Area: 56–63.
- Smith, C.C. 1940. The effect of overgrazing and erosion upon the biota of the mixed grass prairie of Oklahoma. Ecology. 21: 381–397.
- Smith, D.M. 1986. The practice of silviculture. 8th ed. New York: John Wiley and Sons. 527 p.
- Smith, D.M. 1989. Even-aged management: when is it appropriate and what does it reveal about stand development? In: Martin, C. Wayne; Smith, C. Tattersall; Tritton, Louise M., eds. New perspectives on silvicultural management of northern hardwoods: Proceedings of the 1988 symposium on the conflicting consequences of practicing northern hardwood silviculture. 1998 June 9–10; Durham, NH. Gen. Tech. Rep. NE-124. Broomall, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station: 17–25.
- Smith, Kenneth L. 1986. Sawmill: the story of cutting the last great virgin forest east of the Rockies. Fayetteville, AR: University of Arkansas Press. 246 p.
- Smith, R.H.; Lee, R.E., III. 1972. Black turpentine beetle. For. Pest Leaf. 12. Washington, DC: U.S. Department of Agriculture, Forest Service. 8 p.
- Smith, Tim E., ed. 1993. Missouri vegetation management manual. Jefferson City, MO: Missouri Department of Conservation. 148 p.
- Smith, Tim E. 1997. Exotic plant species. In: Missouri Department of Conservation. Conservation policies and guidelines for area and resource management. Jefferson City, MO: Missouri Department of Conservation: VI-74–VI-74c.
- Smith, W. Brad. 1990. Timber resources of Missouri's Northwest Ozarks. Res. Bull. NC-114. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 44 p.
- Soulé, Michael E. 1990. The onslaught of alien species, and other challenges in the coming decades. Conservation Biology. 4(3): 233–239.
- Southern Appalachian Man and the Biosphere (SAMAB). 1996. The southern Appalachian assessment terrestrial technical report. Report 5 of 5. Atlanta, GA: U.S. Department of Agriculture, Forest Service, Southern Region. 288 p.
- Sparks, Jeffrey C.; Masters, Ronald E.; Engle, David M. [and others]. 1998. Effects of late growing-season and late dormant-season prescribed fire on herbaceous vegetation in restored pine-grassland communities. Journal of Vegetation Science. 9: 133–142.
- Spurr, Stephen H.; Barnes, Burton V. 1980. Forest ecology. 3rd ed. New York: John Wiley and Sons. 687 p.
- Stanford, Jack A. 1970. Bobwhite quail. In: W.O. Nagel, ed. Conservation contrasts: three decades of non-political management of wildlife and forests in Missouri. Jefferson City, MO: Missouri Department of Conservation: 44–71.
- Starkey, D.A.; Cost, N.D.; May, D.M. [and others]. 1992. [Plant Pathologist, U.S. Department of Agriculture, Forest Service, State & Private Forestry, Pineville, LA; Project Leader, U.S.

- Department of Agriculture, Forest Service, Southern Forest Experiment Station, Asheville, NC; Research Forester, U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. Forest health monitoring oak decline atlas layer. Atlanta, GA: U.S. Department of Agriculture, Forest Service, Southern Region, State and Private Forestry, Forest Health. 12 p. [Unpublished maps on file at U.S. Forest Service, Southern Region, State and Private Forestry, Forest Health, Asheville, NC.]
- Starkey, D.A.; Oak, S.W.; Ryan, G.W. [and others]. 1989. Evaluation of oak decline areas in the South. Protection Report R8-PR-17. Atlanta, GA: U.S. Department of Agriculture, Forest Service, State and Private Forestry, Southern Region. 36 p.
- Sternitzke, Herbert S. 1960. Arkansas forests. Forest Surv. Rel. 84. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 58 p.
- Sternitzke, Herbert S.; Van Sickle, Charles C. 1968. East Oklahoma forests. Resource Bull. SO-14. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 32 p.
- Stevens, Russell L. 1997. The feral hog in Oklahoma. Ardmore, OK: Samuel Roberts Noble Foundation. 20 p.
- Steyermark, J.A. 1959. Vegetational history of the Ozark forest. Columbia, MO: The University of Missouri. 138 p.
- Stipes, R. Jay; Campana, Richard J., eds. 1981. Compendium of elm diseases. St. Paul, MN: The American Phytopathological Society. 96 p.
- Stout, B.B.; Deschenes, J.M.; Ohmann, L.F. 1975. Multi-species model of a deciduous forest. *Ecology*. 56: 226–231.
- Strausberg, Stephen; Hough, Walter A. 1997. The Ouachita and Ozark-St. Francis National Forests: a history of the lands and the USDA Forest Service tenure. Gen. Tech. Rep. SO-121. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Research Station. 45 p.
- Stringer, J.W.; Kimmerer, T.W.; Overstreet, J.C.; Dunn, J.P. 1989. Oak mortality in eastern Kentucky. *Southern Journal of Applied Forestry*. 13(2): 86–91.
- Strode, Donald D. 1977. Butternut/*Juglans cinerea* L. In: Halls, Lowell K., ed. Southern fruit-producing woody plants used by wildlife. Gen. Tech. Rep. SO-16. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station: 149–150.
- Svejcar, Tony; Christiansen, Tony. 1987. Grazing effects on water relations of Caucasian bluestem. *Journal of Range Management*. 40(1): 15–18.
- Swain, Kenneth M., Sr.; Remion, Michael C. 1974. Direct control methods for the southern pine beetle. USDA Handb. No. 575. Washington, DC: U.S. Department of Agriculture, Forest Service. 15 p.
- Tainter, F.H.; Baker, F.A. 1996. Principles of forest pathology. New York: John Wiley and Sons, Inc.: 571–582, 646–665.
- Tainter, F.H.; Fraedrich, S.W.; Benson, J.D. 1984. The effect of climate on growth, decline, and death of northern red oaks in the western North Carolina Nantahala Mountains. *Castanea*. 49(3): 127–137.
- Tainter, F.H.; Retzlaff, W.A.; Starkey, D.A.; Oak, S.W. 1990. Decline of radial growth in oaks is associated with short-term changes in climate. *European Journal of Forest Pathology*. 20: 95–105.
- Teale, Edwin Way. 1948. Days without time. In: Krutch, Joseph Wood; Eriksson, Paul S. A treasury of birdlore. New York: Joseph Wood Krutch and Paul S. Eriksson: 377–381.
- Terborgh, J. 1989. Where have all the birds gone? Princeton, NJ: Princeton University Press. 207 p.
- Thatcher, R.C.; 1960. Bark beetles affecting southern pines: a review of current knowledge. Occ. Pap. 180. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 25 p.
- Thatcher, R.C.; Barry, P.J. 1982. Southern pine beetle. For. Insect & Dis. Leaf. 49. Washington, DC: U.S. Department of Agriculture, Forest Service. 7 p.
- Thatcher, R.C.; Searcy, J.L.; Coster, J.E.; Hertel, G.D., eds. 1980. The southern pine beetle. Tech. Bull. 1631. Pineville, LA: U.S. Department of Agriculture, Forest Service, Expanded Southern Pine Beetle Research and Applications Program. 266 p.
- Thill, Ronald E. 1984. Deer and cattle diets on Louisiana pine-hardwood sites. *Journal of Wildlife Management*. 48(3): 788–798.
- Thompson, F. R., III. 1994. Temporal and spatial patterns of breeding brown-headed cowbirds in the Midwestern United States. *The Auk*. 111: 979–990.
- Thompson, F. R., III. 1995. Management of midwestern landscapes for the conservation of neotropical migratory birds. Gen. Tech. Rep. NC-187. St. Paul, MN: U.S. Department of Agriculture, Forest Service. 208 p.
- Thompson, F. R., III.; Lewis, S.J.; Green, J.; Ewert, D. 1993. Status of neotropical landbirds in the Midwest: Identifying species of management concern. In: Finch, D. M.; Stangel, P. W., eds. Status and management of neotropical migratory

- birds. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station: 145–158.
- Thompson, F. R., III.; Probst, J.R.; Raphael, M.G. 1995. Impacts of silviculture: overview and management recommendations. In: Martin, T. E.; Finch, D. M., eds. Ecology and management of neotropical migratory birds. New York: Oxford University Press: 201–219.
- Thompson, F. R., III.; Robinson, S.K.; Donovan, T.M. [and others]. [In press]. Biogeographic, landscape, and local factors affecting cowbird abundance and host parasitism levels. In: Cook, T.; Robinson, S.K.; Rothstein, S.I.; Sealy, S.G., eds. The ecology and management of cowbirds. Austin, TX: University of Texas Press.
- Thompson, F. R., III.; Robinson, S.K.; Whitehead, D.R.; Brawn, J.D. 1996. Management of central hardwood landscapes for the conservation of migratory birds. In: Thompson, F. R., III, ed. Management of midwestern landscapes for the conservation of neotropical migratory birds. Gen. Tech. Rep. NC-187. St. Paul, MN: U.S. Department of Agriculture, Forest Service: 117–143.
- Thornbury, William D. 1965. Regional geomorphology of the United States. New York: John Wiley and Sons. 609 pages.
- Tigner, Tim. 1992. Gypsy moth impact on Virginia's hardwood forests and forest industry. Charlottesville, VA: Virginia Department of Forestry. 35 p.
- Trimble, G.R., Jr. 1970. 20 years of intensive uneven-aged management: effect on growth, yield, and species composition in two hardwood stands in West Virginia. Res. Pap. NE-154. Upper Darby, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station. 12 p.
- Trimble, G.R., Jr. 1973. The regeneration of Central Appalachian hardwoods with emphasis on the effects of site quality and harvesting practice. Res. Pap. NE-282. Upper Darby, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station. 14 p.
- Triplehorn, Charles A. 1955. The Asiatic oak weevil in Delaware. *Journal of Economic Entomology*. 48(3): 289–293.
- Tryon, E.H.; Powell, D.S. 1984. Root ages of advance hardwood reproduction. *Forest Ecology Management*. 8: 293–298.
- Tyrrell, Lucy E.; Nowacki, Gregory J.; Crow, Thomas R. [and others]. 1998. Information about old growth for selected forest type groups in the Eastern United States. Gen. Tech. Rep. NC-197. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 507 p.
- U.S. Department of Agriculture, Forest Service (USDA FS). 1986. Land and resource management plan, Mark Twain National Forest. Milwaukee, WI: U.S. Department of Agriculture, Forest Service, Eastern Region. 252 p. + appendices.
- U.S. Department of Agriculture, Forest Service (USDA FS). 1987. Final environmental impact statement for the suppression of the southern pine beetle—southern region. Manage. Bull. R8-MB2. Atlanta, GA: U.S. Department of Agriculture, Forest Service, Southern Region. 1,627 p. Vol. I–III.
- U.S. Department of Agriculture, Forest Service (USDA FS). 1989. Interim management guide for Ozark chinquapin (*Castanea pumila* var. *ozarkensis*). U.S. Department of Agriculture, Forest Service, Ozark National Forest. 32 p. [Unpublished document on file at the Supervisor's office, U.S. Department of Agriculture, Forest Service, Ozark National Forest, Russellville, AR.]
- U.S. Department of Agriculture, Forest Service (USDA FS). 1990. Record of decision, final environmental impact statement, vegetation management in the Ozark/Ouachita Mountains, March 5, 1990. Atlanta, GA: U.S. Department of Agriculture, Forest Service, Southern Region. 17 p.
- U.S. Department of Agriculture, Forest Service (USDA FS). 1995a. Course to the future: positioning fire and aviation management. Washington, DC: U.S. Department of Agriculture, Forest Service, Fire and Aviation Management. 30 p.
- U.S. Department of Agriculture, Forest Service (USDA FS). 1995b. Gypsy moth management in the United States: a cooperative approach: final Environmental Impact Statement. Washington, DC: U.S. Department of Agriculture, Forest Service. 704 p.
- U.S. Department of Agriculture, Forest Service (USDA FS). 1996. Environmental impact statement for the renewal of the shortleaf pine/bluestem grass ecosystem and recovery of the red-cockaded woodpecker. Hot Springs, AR: U.S. Department of Agriculture, Forest Service. 48 p.
- U.S. Department of Agriculture, Forest Service (USDA FS). [Accessed October 1997]. Fire effects information. <<http://www.fs.fed.us/database/feis>> Missoula, MT: U.S. Department of Agriculture, Forest Service. [Various pagination].
- U.S. Department of Agriculture, Forest Service (USDA FS). 1999a. Ozark-Ouachita Highlands Assessment: aquatic conditions. Gen. Tech. Rep. SRS-33. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Research Station. 318 p.
- U.S. Department of Agriculture, Forest Service (USDA FS). 1999b. Ozark-Ouachita Highlands Assessment: social and

- economic conditions. Gen. Tech. Rep. SRS-34. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Research Station. 300 p.
- U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station (USDA FS SFES). 1956. Forests of east Oklahoma, 1955–56. Forest Surv. Rel. 79. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 34 p.
- U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station (USDA FS SFES). 1979. Forest statistics for Arkansas counties. Resour. Bull. SO-76. New Orleans, LA: Southern Forest Experiment Station. 77 p.
- U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station (USDA FS SFES). 1992. Forest maps of the United States. 1993 RPA Program, compact disc. Starkville, MS: U.S. Department of Agriculture, Forest Service, Southern Research Station, Forest Inventory and Analysis. [Not paged].
- U.S. Department of Agriculture, Forest Service, Southern Research Station (USDA FS SRS). [Accessed February 1997]. Forest inventory and analysis data base retrieval system. <<http://www.srsfia.usfs.msstate.edu/scripts/ew.htm>>. Starkville, MS: U.S. Department of Agriculture, Forest Service, Southern Research Station, Forest Inventory and Analysis.
- U.S. Department of Agriculture, Natural Resources Conservation Service (USDA NRCS). 1997. National Resource Inventory (NRI). Hot Springs, AR: Ouachita National Forest. 16 p. [Unpublished data summary on file at the Forest Supervisor's Office, U.S. Department of Agriculture, Forest Service, Ouachita National Forest, Hot Springs, AR.]
- Van Lear, D.; Watt, J.M. 1992. The role of fire in oak regeneration. In: Loftis, D.L.; McGee, C.E., eds. Oak regeneration: serious problems and practical recommendations. Gen. Tech. Rept. SE-84. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station: 66–78.
- Vasilevsky, Alexander; Essex, Burton L. 1974. Timber resources of Missouri's northwestern Ozarks. Resour. Bull. NC-22. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 59 p.
- Venator, C.R. 1985. Survival of shortleaf pine (*Pinus echinata* Mill.) seedlings as influenced by nursery handling and storage. Tree Planters' Notes. 36(4): 17–19.
- Walker, W.D. 1992. Historical perspectives on regeneration in the Ouachita and Ozark Mountains—the Ouachita National Forest. In: Brissette, J.C.; Barnett, J.P., comps. Proceedings of the shortleaf pine regeneration workshop; 1991 October 29–31; Little Rock, AR. Gen. Tech. Rep. SO-90. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station: 12–17.
- Ward, W.W. 1966. Oak-hardwood reproduction in central Pennsylvania. Journal of Forestry. 64(11): 744–749.
- Wargo, P.M.; Houston, D.R.; LaMadeleine, L.A. 1983. Oak decline [Leaflet]. For. Insect & Dis. Leaf. 165. Washington, DC: U.S. Department of Agriculture, Forest Service. 8 p.
- Weakley, Alan; Patterson, K.D.; Landaal, S., Pyne, M. 1996. International classification of ecological communities: terrestrial vegetation of the Southeastern United States. Chapel Hill, NC: The Nature Conservancy, Southeast Regional Office, Southern Conservation Science Department, Community Ecology Group. 374 p.
- Weakley, Alan; Patterson, K.D.; Landaal, Pyne, M. 1997. International classification of ecological communities: terrestrial vegetation of the Southeastern United States. Ozark Mountains/Ouachita Mountains—draft subset (alliance level). Chapel Hill, NC: The Nature Conservancy, Southeast Regional Office, Southern Conservation Science Department, Community Ecology Group. 113 p.
- Weaver, Dwight H. 1992. The wilderness underground, caves of the Ozark plateau. Columbia, MO: University of Missouri Press. 113 p.
- Weitzman, S.; Trimble, G.R., Jr. 1957. Some natural factors that govern the management of oaks. Stn. Pap. No. 88. Upper Darby, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station. 40 p.
- Wells, O.O. 1969. Results of the southwide seed source study through 1968–1969. Proceedings of the tenth southern conference on forest tree improvement. 1969 June 17–19; Houston, TX. College Station, TX, Texas Forest Service: 117–129.
- Wescott, C. 1964. The gardener's bug book. New York: Doubleday. 625 p.
- White, P.H. 1979. Pattern, process, and natural disturbance in vegetation. The Botanical Review. 45: 229–299.
- Williams, G.W. 1994. [Sociologist and Social Historian, USDA Forest Service, Pacific Northwest Region, Portland, OR] References on the American Indian use of fire in ecosystems. Portland, OR: United States Department of Agriculture, Forest Service. 24 p. [Unpublished report on file at U.S. Department of Agriculture, Forest Service, Portland, OR.]

- Williams, Jerry. 1993. Fire related considerations and strategies in support of ecosystem management. Washington, DC: U.S. Department of Agriculture, Forest Service, Fire and Aviation Management [Staff paper]. 30 p.
- Williams, Ted. 1994. Invasion of the aliens. Audubon. (Sept.–Oct.): 24–32.
- Wilson, Christopher W.; Masters, Ronald E.; Bukenhofer, George A. 1995. Breeding bird response to pine-grassland community restoration for red-cockaded woodpeckers. *The Journal of Wildlife Management*. 59(1): 56–67.
- Wilson, Louis F. 1971. Walkingstick. *Forest Pest Leaf*. 82. Washington, DC: U.S. Department of Agriculture, Forest Service. 4 p.
- Winkel, Von K.; Roundy, Bruce A. 1991. Effects of cattle trampling and mechanical seedbed preparation on grass seedling emergence. *Journal of Range Management*. 44(2): 176–178.
- Winters, R.K.; Roberts, E.V. 1948a. Forest resources of the eastern Ozark region in Missouri. *Forest Surv. Rel. No. 1*. Columbus, OH: U.S. Department of Agriculture, Forest Service, Central States Forest Experiment Station. 21 p.
- Winters, R.K.; Roberts, E.V. 1948b. Forest resources of the northwest Ozark region in Missouri. *Forest Surv. Rel. No. 3*. Columbus, OH: U.S. Department of Agriculture, Forest Service, Central States Forest Experiment Station. 19 p.
- Winters, R.K.; Roberts, E.V. 1948c. Forest resources of the southwestern Ozark region in Missouri. *Forest Surv. Rel. No. 2*. Columbus, OH: U.S. Department of Agriculture, Forest Service, Central States Forest Experiment Station. 21 p.
- Wittwer, R.F.; Dougherty, P.M.; Cosby, D. 1986. Effects of ripping and herbicide site preparation treatments on loblolly pine seedling growth and survival. *Southern Journal of Applied Forestry*. 10(4): 253–257.
- Wittwer, R.F.; Lynch, T.B.; Huebschmann, M.M. 1996. Thinning improves growth of crop trees in natural shortleaf pine stands. *Southern Journal of Applied Forestry*. 20(4): 182–187.
- Wojcik, D.P.; Buren, W.F.; Grissell, E.E.; Cardysle, T. 1976. The fire ants (*Solenopsis*) of Florida (*Hymenoptera: Formicidae*). *Entomolog. Circ. No. 173*. Florida Department of Agriculture and Consumer Services. 4 p.
- Woods, E.D.; Wittwer, R.F.; Dougherty, P.M. [and others]. 1988. Influence of site factors on growth of loblolly and shortleaf pine in Oklahoma. Research Report No. P-900. Stillwater, OK: Oklahoma State University, Division of Agriculture, Agricultural Experiment Station. 25 p.
- Yeiser, J.L. 1986. Tree injection for early pine seedling release in the Ozark Mountains of Arkansas. *Southern Journal of Applied Forestry*. 10(4): 249–251.
- Yeiser, J.L.; Barnett, J.P. 1991. Growth and physiological response of four shortleaf pine families to herbicidal control of herbaceous competition. *Southern Journal of Applied Forestry*. 15 (4): 199–208.
- Yeiser, J.L.; Cobb, S.W. 1988. Herbicides for releasing pine seedlings in the Arkansas Ozarks. *Bull. No. 911*. Fayetteville, AR: University of Arkansas, Arkansas Agricultural Experiment Station. 13 p.
- Yeiser, J.L.; Sundell, E.S.; Boyd, J.W. 1987. Preplant and postplant treatments for weed control in newly planted pine. *Bull. No. 902*. Fayetteville, AR: University of Arkansas, Arkansas Agricultural Experiment Station. 10 p.
- Yocom, H.A. 1971. Shortleaf pine seed dispersal. *Journal of Forestry*. 66(5): 422.
- Yocom, H.A. 1972. Burning to reduce understory hardwoods in the Arkansas mountains. *Res. Note SO-145*. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 3 p.
- Yocom, H.A.; Lawson, E.R. 1977. Tree percent from naturally regenerated shortleaf pine. *Southern Journal of Applied Forestry*. 1(2): 10–11.
- Young, J.A.; Young, C.G. 1992. *Seeds of woody plants in North America*. 2nd ed. Portland, OR: Dioscoridies Press. 407 p.

U.S. Department of Agriculture, Forest Service. 1999. Ozark-Ouachita Highlands Assessment: terrestrial vegetation and wildlife. Report 5 of 5. Gen. Tech. Rep. SRS-35. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Research Station. 201 p.

This publication provides citizens, private and public organizations, scientists, and others with information about terrestrial animals, plants, and biological communities in and near the national forests in the Ozark-Ouachita Highlands: the Mark Twain in Missouri, the Ouachita in Arkansas and Oklahoma, and the Ozark-St. Francis National Forests in Arkansas. The document examines the status and trends of vegetation, plant and animal populations, forest management, and biological threats to forest resources in the Highlands.

Keywords: Biological threats, ecological classification, forest management, plant and animal populations, silviculture, vegetation cover.